

# Coordinate Descent

Ryan Tibshirani  
Convex Optimization 10-725

## Last time: numerical linear algebra primer

In  $\mathbb{R}^n$ , rough **flop counts** for basic operations are as follows

- Vector-vector operations:  $n$  flops
- Matrix-vector multiplication:  $n^2$  flops
- Matrix-matrix multiplication:  $n^3$  flops
- Linear system solve:  $n^3$  flops

Operations with **banded or sparse matrices** are much cheaper

Two classes of approaches for linear system solvers:

- **Direct**: QR decomposition, Cholesky decomposition
- **Indirect**: Jacobi, Gauss-Seidl, gradient descent, conjugate gradient

Rough rule of thumb: if problem fits easily in memory, go direct, else go indirect

# Outline

Today:

- Coordinate descent
- Examples
- Implementation tricks
- Graphical lasso
- Screening rules

## Coordinatewise optimality


We've seen some pretty sophisticated methods thus far

We now focus on a very simple technique that can be surprisingly efficient, scalable: **coordinate descent**, or more appropriately called coordinatewise minimization

Q: Given convex, differentiable  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , if we are at a point  $x$  such that  $f(x)$  is minimized along each coordinate axis, then *have we found a global minimizer?*

That is, does  $f(x + \delta e_i) \geq f(x)$  for all  $\delta, i \Rightarrow f(x) = \min_z f(z)$ ?

(Here  $e_i = (0, \dots, 1, \dots, 0) \in \mathbb{R}^n$  is the  $i$ th standard basis vector)





A: Yes! Proof:

$$0 = \nabla f(x) = \left( \frac{\partial f}{\partial x_1}(x), \dots, \frac{\partial f}{\partial x_n}(x) \right)$$

---



Q: Same question, but now for  $f$  convex, and not differentiable?



A: No! Look at the above counterexample

---

Q: Same, now  $f(x) = g(x) + \sum_{i=1}^n h_i(x_i)$ , with  $g$  convex, smooth, and each  $h_i$  convex? (Here the nonsmooth part is called **separable**)



A: Yes! Proof: using convexity of  $g$  and subgradient optimality

$$\begin{aligned}
 f(y) - f(x) &= g(x) - g(y) + \sum_{i=1}^n [h_i(y_i) - h_i(x_i)] \\
 &\geq \sum_{i=1}^n \underbrace{[\nabla_i g(x)(y_i - x_i) + h_i(y_i) - h_i(x_i)]}_{\geq 0} \geq 0
 \end{aligned}$$

# Coordinate descent

This suggests that for the problem

$$\min_x f(x)$$

where  $f(x) = g(x) + \sum_{i=1}^n h_i(x_i)$ , with  $g$  convex and differentiable and each  $h_i$  convex, we can use **coordinate descent**: let  $x^{(0)} \in \mathbb{R}^n$ , and repeat

$$x_i^{(k)} = \operatorname{argmin}_{x_i} f(x_1^{(k)}, \dots, x_{i-1}^{(k)}, x_i, x_{i+1}^{(k-1)}, \dots, x_n^{(k-1)}),$$
$$i = 1, \dots, n$$

for  $k = 1, 2, 3, \dots$

Important note: we always use **most recent information** possible

Tseng (2001) showed that for such  $f$  (provided  $f$  is continuous on compact set  $\{x : f(x) \leq f(x^{(0)})\}$  and  $f$  attains its minimum), any limit point of  $x^{(k)}$ ,  $k = 1, 2, 3, \dots$  is a minimizer of  $f$ <sup>1</sup>

Notes:

- Order of cycle through coordinates is arbitrary, can use any permutation of  $\{1, 2, \dots, n\}$
- Can everywhere replace individual coordinates with blocks of coordinates
- “One-at-a-time” update scheme is critical, and “all-at-once” scheme **does not** necessarily converge
- The analogy for solving linear systems: Gauss-Seidel versus Jacobi method

---

<sup>1</sup>Using basic real analysis, we know  $x^{(k)}$  has subsequence converging to  $x^*$  (Bolzano-Weierstrass), and  $f(x^{(k)})$  converges to  $f^*$  (monotone convergence)

## Example: linear regression

Given  $y \in \mathbb{R}^n$ , and  $X \in \mathbb{R}^{n \times p}$  with columns  $X_1, \dots, X_p$ , consider the **linear regression** problem:

$$\min_{\beta} \frac{1}{2} \|y - X\beta\|_2^2$$

Minimizing over  $\beta_i$ , with all  $\beta_j$ ,  $j \neq i$  fixed:


$$0 = \nabla_i f(\beta) = X_i^T (X\beta - y) = X_i^T (X_i\beta_i + X_{-i}\beta_{-i} - y)$$

i.e., we take

$$\beta_i = \frac{X_i^T (y - X_{-i}\beta_{-i})}{X_i^T X_i}$$

Coordinate descent repeats this update for  $i = 1, 2, \dots, p, 1, 2, \dots$ . Note that this is exactly **Gauss-Seidl** for the system  $X^T X\beta = X^T y$

Coordinate descent vs gradient descent for linear regression: 100 random instances with  $n = 100$ ,  $p = 20$



Is it fair to compare 1 cycle of coordinate descent to 1 iteration of gradient descent? Yes, if we're clever

- Gradient descent:  $\beta \leftarrow \beta + tX^T(y - X\beta)$ , costs  $O(np)$  flops
- Coordinate descent, one coordinate update:

$$\beta_i \leftarrow \frac{X_i^T(y - X_{-i}\beta_{-i})}{X_i^T X_i} = \frac{X_i^T r}{\|X_i\|_2^2} + \beta_i$$

where  $r = y - X\beta$

- Each coordinate costs  $O(n)$  flops:  $O(n)$  to update  $r$ ,  $O(n)$  to compute  $X_i^T r$
- One cycle of coordinate descent costs  $O(np)$  operations, **same as gradient descent**

## Example: lasso regression

Consider the **lasso** problem:

$$\min_{\beta} \frac{1}{2} \|y - X\beta\|_2^2 + \lambda \|\beta\|_1$$

Note that nonsmooth part here is separable:  $\|\beta\|_1 = \sum_{i=1}^p |\beta_i|$ .  
Minimizing over  $\beta_i$ , with  $\beta_j$ ,  $j \neq i$  fixed:


$$0 = X_i^T X_i \beta_i + X_i^T (X_{-i} \beta_{-i} - y) + \lambda s_i$$

where  $s_i \in \partial |\beta_i|$ . Solution is simply given by soft-thresholding

$$\beta_i = S_{\lambda / \|X_i\|_2^2} \left( \frac{X_i^T (y - X_{-i} \beta_{-i})}{X_i^T X_i} \right)$$

Repeat this for  $i = 1, 2, \dots, p, 1, 2, \dots$

Coordinate descent vs proximal gradient for lasso regression: 100 random instances with  $n = 200$ ,  $p = 50$  (all methods cost  $O(np)$  per iter)



## Example: box-constrained QP

Given  $b \in \mathbb{R}^n$ ,  $Q \in \mathbb{S}_+^n$ , consider a **box-constrained QP**:

$$\min_x \frac{1}{2} x^T Q x + b^T x \quad \text{subject to} \quad l \leq x \leq u$$

Fits into our framework, as  $I\{l \leq x \leq u\} = \sum_{i=1}^n I\{l_i \leq x_i \leq u_i\}$

Minimizing over  $x_i$  with all  $x_j$ ,  $j \neq i$  fixed: same basic steps give

$$x_i = T_{[l_i, u_i]} \left( \frac{b_i - \sum_{j \neq i} Q_{ij} x_j}{Q_{ii}} \right)$$

where  $T_{[l_i, u_i]}$  is the truncation (projection) operator onto  $[l_i, u_i]$ :

$$T_{[l_i, u_i]}(z) = \begin{cases} u_i & \text{if } z > u_i \\ z & \text{if } l_i \leq z \leq u_i \\ l_i & \text{if } z < l_i \end{cases}$$

## Example: support vector machines

A coordinate descent strategy can be applied to the **SVM dual**:

$$\min_{\alpha} \frac{1}{2} \alpha^T \tilde{X} \tilde{X}^T \alpha - 1^T \alpha \quad \text{subject to} \quad 0 \leq \alpha \leq C \mathbf{1}, \quad \alpha^T y = 0$$

**Sequential minimal optimization** or SMO (Platt 1998) is basically blockwise coordinate descent in blocks of 2. Instead of cycling, it chooses the next block greedily

Recall the complementary slackness conditions

$$\alpha_i (1 - \xi_i - (\tilde{X} \beta)_i - y_i \beta_0) = 0, \quad i = 1, \dots, n \quad (1)$$


$$(C - \alpha_i) \xi_i = 0, \quad i = 1, \dots, n \quad (2)$$

where  $\beta, \beta_0, \xi$  are the primal coefficients, intercept, and slacks. Recall that  $\beta = \tilde{X}^T \alpha$ ,  $\beta_0$  is computed from (1) using any  $i$  such that  $0 < \alpha_i < C$ , and  $\xi$  is computed from (1), (2)

SMO repeats the following two steps:

- Choose  $\alpha_i, \alpha_j$  that violate complementary slackness, greedily (using heuristics)
- Minimize over  $\alpha_i, \alpha_j$  exactly, keeping all other variables fixed

Using equality constraint, reduces to minimizing univariate quadratic over an interval (From Platt 1998)



Note this does not meet separability assumptions for convergence from Tseng (2001), and a different treatment is required

Many further developments on coordinate descent for SVMs have been made; e.g., a recent one is Hsieh et al. (2008)

# Coordinate descent in statistics and ML

History in statistics/ML:

- Idea appeared in Fu (1998), and then again in Daubechies et al. (2004), but was inexplicably ignored
- Later, three papers in 2007, especially Friedman et al. (2007), really sparked interest in statistics and ML communities

Why is it used?

- Very simple and easy to implement
- Careful implementations can achieve state-of-the-art
- Scalable, e.g., don't need to keep full data in memory

Examples: lasso regression, lasso GLMs (under proximal Newton), SVMs, group lasso, graphical lasso (applied to the dual), additive modeling, matrix completion, regression with nonconvex penalties

## Pathwise coordinate descent for lasso

Structure for pathwise coordinate descent, Friedman et al. (2009):

Outer loop (**pathwise** strategy):

- Compute the solution over a sequence  $\lambda_1 > \lambda_2 > \dots > \lambda_r$  of tuning parameter values
- For tuning parameter value  $\lambda_k$ , initialize coordinate descent algorithm at the computed solution for  $\lambda_{k+1}$  (warm start)

Inner loop (**active set** strategy):

- Perform one coordinate cycle (or small number of cycles), and record active set  $A$  of coefficients that are nonzero
- Cycle over only the coefficients in  $A$  until convergence
- Check KKT conditions over all coefficients; if not all satisfied, add offending coefficients to  $A$ , go back one step

## Notes:

- Even when the solution is desired at only one  $\lambda$ , the pathwise strategy (solving over  $\lambda_1 > \dots > \lambda_r = \lambda$ ) is typically much more efficient than directly performing coordinate descent at  $\lambda$
- Active set strategy takes advantage of sparsity; e.g., for large problems, coordinate descent for lasso is much faster than it is for ridge regression
- With these strategies in place (and a few more clever tricks), coordinate descent can be competitive with fastest algorithms for  $\ell_1$  penalized minimization problems
- Fortran implementation [glmnet](#), linked to R, MATLAB, etc.

## Coordinate gradient descent

For a smooth function  $f$ , the iterations

$$x_i^{(k)} = x_i^{(k-1)} - t_{ki} \cdot \nabla_i f(x_1^{(k)}, \dots, x_{i-1}^{(k)}, x_i, x_{i+1}^{(k-1)}, \dots, x_n^{(k-1)}), \\ i = 1, \dots, n$$

for  $k = 1, 2, 3, \dots$  are called **coordinate gradient descent**, and when  $f = g + h$ , with  $g$  smooth and  $h = \sum_{i=1}^n h_i$ , the iterations

$$x_i^{(k)} = \text{PROX}_{h_i, t_{ki}} \left( x_i^{(k-1)} - t_{ki} \cdot \nabla_i g(x_1^{(k)}, \dots, x_i^{(k-1)}, \dots, x_n^{(k-1)}) \right), \\ i = 1, \dots, n$$

for  $k = 1, 2, 3, \dots$  are called **coordinate proximal gradient descent**

When  $g$  is quadratic, (proximal) coordinate gradient descent is the same as coordinate descent under proper step sizes

## Convergence analyses

Theory for coordinate descent moves quickly. Each combination of the following cases has (probably) been analyzed:

- Coordinate descent or (proximal) coordinate gradient descent?
- Cyclic rule, permuted cyclic, or greedy rule, randomized rule?

Roughly speaking, results are similar to those for proximal gradient descent: under standard conditions, get standard rates

But **constants differ** and this matters! Much recent work is focused on improving them

Also, it is generally believe that coordinate descent should perform better than first-order methods (when implementable)

Some references are Beck and Tetruashvili (2013), Wright (2015), Sun and Hong (2015), Li et al. (2016)

# Graphical lasso

Consider  $X \in \mathbb{R}^{n \times p}$ , with rows  $x_i \sim N(0, \Sigma)$ ,  $i = 1, \dots, n$ , drawn independently. Suppose  $\Sigma$  is unknown. It is often reasonable (for large  $p$ ) to seek a sparse estimate of  $\Sigma^{-1}$

Why? For  $z \sim N(0, \Sigma)$ , normality theory tells us

$$\Sigma_{ij}^{-1} = 0 \iff z_i, z_j \text{ conditionally independent given } z_\ell, \ell \neq i, j$$

**Graphical lasso** (Banerjee et al. 2007, Friedman et al. 2007):

$$\min_{\Theta \in \mathbb{S}_+^p} -\log \det \Theta + \text{tr}(S\Theta) + \lambda \|\Theta\|_1$$

where  $S = X^T X/n$ , and  $\|\Theta\|_1 = \sum_{i,j=1}^p |\Theta_{ij}|$ . Observe that this is a convex problem. Solution  $\hat{\Theta}$  serves as estimate for  $\Sigma^{-1}$

## Gllasso algorithm

Graphical lasso KKT conditions (stationarity):

$$-\Theta^{-1} + S + \lambda\Gamma = 0$$

where  $\Gamma_{ij} \in \partial|\Theta_{ij}|$ . Let  $W = \Theta^{-1}$ . Note  $W_{ii} = S_{ii} + \lambda$ , because  $\Theta_{ii} > 0$  at solution. Now partition:

$$\begin{array}{c} W = \\ \left[ \begin{array}{cc} W_{11} & w_{12} \\ w_{21} & w_{22} \end{array} \right] \end{array} \quad \begin{array}{c} \Theta = \\ \left[ \begin{array}{cc} \Theta_{11} & \theta_{12} \\ \theta_{21} & \theta_{22} \end{array} \right] \end{array} \quad \begin{array}{c} S = \\ \left[ \begin{array}{cc} S_{11} & s_{12} \\ s_{21} & s_{22} \end{array} \right] \end{array} \quad \begin{array}{c} \Gamma = \\ \left[ \begin{array}{cc} \Gamma_{11} & \gamma_{12} \\ \gamma_{21} & \gamma_{22} \end{array} \right] \end{array}$$

where  $W_{11} \in \mathbb{R}^{(p-1) \times (p-1)}$ ,  $w_{12} \in \mathbb{R}^{(p-1) \times 1}$ , and  $w_{21} \in \mathbb{R}^{1 \times (p-1)}$ ,  $w_{22} \in \mathbb{R}$ ; same with others

**Gllasso algorithm** (Friedman et al., 2007): solve for  $w_{12}$  (recall that  $w_{22}$  is known), with all other columns fixed; then solve for second-to-last column, etc., and cycle around until convergence

## Screening rules

In some problems, **screening rules** can be used in combination with coordinate descent to further wittle down the active set. Screening rules themselves have amassed a huge literature recently

Originated with El Ghaoui et al. (2010), SAFE rule for the lasso:

$$|X_i^T y| < \lambda - \|X_i\|_2 \|y\|_2 \frac{\lambda_{\max} - \lambda}{\lambda_{\max}} \Rightarrow \hat{\beta}_i = 0, \quad \text{all } i = 1, \dots, p$$

where  $\lambda_{\max} = \|X^T y\|_{\infty}$  (the smallest value of  $\lambda$  such that  $\hat{\beta} = 0$ )

Note: this is **not an if and only if** statement! But it does give us a way of eliminating features apriori, without solving the lasso

(There have been many advances in screening rules for the lasso ... but this was the first, and one of the simplest)

## SAFE rule derivation

Why is the SAFE rule true? Construction comes from **lasso dual**:

$$\max_u g(u) \quad \text{subject to} \quad \|X^T u\|_\infty \leq \lambda$$

where  $g(u) = \frac{1}{2}\|y\|_2^2 - \frac{1}{2}\|y - u\|_2^2$ . Suppose that  $u_0$  is dual feasible (e.g., take  $u_0 = y \cdot \lambda / \lambda_{\max}$ ). Then  $\gamma = g(u_0)$  is a lower bound on the dual optimal value, so dual problem is equivalent to


$$\max_u g(u) \quad \text{subject to} \quad \|X^T u\|_\infty \leq \lambda, \quad g(u) \geq \gamma$$

Now let  $m_i = \max_u |X_i^T u|$  subject to  $g(u) \geq \gamma$ , for  $i = 1, \dots, p$ .

Then

$$m_i < \lambda \quad \Rightarrow \quad |X_i^T \hat{u}| < \lambda \quad \Rightarrow \quad \hat{\beta}_i = 0, \quad i = 1, \dots, p$$

The last implication comes from the KKT conditions



(From El Ghaoui et al. 2010)

Another dual argument shows that

$$\begin{aligned} & \max_u X_i^T u \quad \text{subject to} \quad g(u) \geq \gamma \\ &= \min_{\mu > 0} -\gamma\mu + \frac{1}{\mu} \|\mu y - X_i\|_2^2 \\ &= \|X_i\|_2 \sqrt{\|y\|_2^2 - 2\gamma} - X_i^T y \end{aligned}$$

where the last equality comes from direct calculation

Thus  $m_i$  is given the maximum of the above quantity over  $\pm X_i$ ,

$$m_i = \|X_i\|_2 \sqrt{\|y\|_2^2 - 2\gamma} + |X_i^T y|, \quad i = 1, \dots, p$$

Lastly, substitute  $\gamma = g(y \cdot \lambda / \lambda_{\max})$ . Then  $m_i < \lambda$  is precisely the SAFE rule given on previous slide

# References

Early coordinate descent references in optimization:

- D. Bertsekas and J. Tsitsiklis (1989), “Parallel and distributed computation: numerical methods”, Chapter 3
- Z. Luo and P. Tseng (1992), “On the convergence of the coordinate descent method for convex differentiable minimization”
- J. Ortega and W. Rheinboldt (1970), “Iterative solution of nonlinear equations in several variables”
- P. Tseng (2001), “Convergence of a block coordinate descent method for nondifferentiable minimization”
- J. Warga (1969), “Minimizing certain convex functions”

## Early coordinate descent references in statistics and ML:

- I. Daubechies and M. Defrise and C. De Mol (2004), “An iterative thresholding algorithm for linear inverse problems with a sparsity constraint”
- J. Friedman and T. Hastie and H. Hoefling and R. Tibshirani (2007), “Pathwise coordinate optimization”
- W. Fu (1998), “Penalized regressions: the bridge versus the lasso”
- T. Wu and K. Lange (2008), “Coordinate descent algorithms for lasso penalized regression”
- A. van der Kooij (2007), “Prediction accuracy and stability of regression with optimal scaling transformations”

## Applications of coordinate descent:

- O. Banerjee and L. Ghaoui and A. d'Aspremont (2007), "Model selection through sparse maximum likelihood estimation"
- J. Friedman and T. Hastie and R. Tibshirani (2007), "Sparse inverse covariance estimation with the graphical lasso"
- J. Friedman and T. Hastie and R. Tibshirani (2009), "Regularization paths for generalized linear models via coordinate descent"
- C.J. Hsieh and K.W. Chang and C.J. Lin and S. Keerthi and S. Sundararajan (2008), "A dual coordinate descent method for large-scale linear SVM"
- R. Mazumder and J. Friedman and T. Hastie (2011), "SparseNet: coordinate descent with non-convex penalties"
- J. Platt (1998), "Sequential minimal optimization: a fast algorithm for training support vector machines"

## Recent theory for coordinate descent:

- A. Beck and L. Tetruashvili (2013), “On the convergence of block coordinate descent type methods”
- X. Li, T. Zhao, R. Arora, H. Liu, M. Hong (2016) “An improved convergence analysis of cyclic block coordinate descent-type methods for strongly convex minimization”
- Y. Nesterov (2010), “Efficiency of coordinate descent methods on huge-scale optimization problems”
- J. Nutini, M. Schmidt, I. Laradji, M. Friedlander, H. Koepke (2015), “Coordinate descent converges faster with the Gauss-Southwell rule than random selection”
- P. Richtarik and M. Takac (2011), “Iteration complexity of randomized block-coordinate descent methods for minimizing a composite function”
- R. Sun and M. Hong (2015), “Improved iteration complexity bounds of cyclic block coordinate descent for convex problems”
- S. Wright (2015), “Coordinate descent algorithms”

## Screening rules and graphical lasso:

- L. El Ghaoui and V. Viallon and T. Rabbani (2010), “Safe feature elimination in sparse supervised learning”
- R. Tibshirani, J. Bien, J. Friedman, T. Hastie, N. Simon, J. Taylor, and R. J. Tibshirani (2011), “Strong rules for discarding predictors in lasso-type problems”
- R. Mazumder and T. Hastie (2011), “The graphical lasso: new insights and alternatives”
- R. Mazumder and T. Hastie (2011), “Exact covariance thresholding into connected components for large-scale graphical lasso”
- J. Wang, P. Wonka, and J. Ye (2015), “Lasso screening rules via dual polytope projection”
- D. Witten and J. Friedman and N. Simon (2011), “New insights and faster computations for the graphical lasso”