

Combinatorial Bounds for Conflict-free Coloring on Open Neighborhoods

Sriram Bhyravarapu and Subrahmanyam Kalyanasundaram

Department of Computer Science and Engineering, IIT Hyderabad
{cs16resch11001, subruk}@iith.ac.in

Abstract. In an undirected graph G , a conflict-free coloring with respect to open neighborhoods (denoted by CFON coloring) is an assignment of colors to the vertices such that every vertex has a uniquely colored vertex in its open neighborhood. The minimum number of colors required for a CFON coloring of G is the CFON chromatic number of G , denoted by $\chi_{ON}(G)$.

The decision problem that asks whether $\chi_{ON}(G) \leq k$ is NP-complete. Structural as well as algorithmic aspects of this problem have been well studied. We obtain the following results for $\chi_{ON}(G)$:

- Bodlaender, Kolay and Pieterse [WADS 2019] showed the upper bound $\chi_{ON}(G) \leq \text{fvs}(G) + 3$, where $\text{fvs}(G)$ denotes the size of a minimum feedback vertex set of G . We show the improved bound of $\chi_{ON}(G) \leq \text{fvs}(G) + 2$, which is tight, thereby answering an open question in the above paper.
- We study the relation between $\chi_{ON}(G)$ and the pathwidth of the graph G , denoted $\text{pw}(G)$. The above paper from WADS 2019 showed the upper bound $\chi_{ON}(G) \leq 2\text{tw}(G) + 1$ where $\text{tw}(G)$ stands for the treewidth of G . This implies an upper bound of $\chi_{ON}(G) \leq 2\text{pw}(G) + 1$. We show an improved bound of $\chi_{ON}(G) \leq \lfloor \frac{5}{3}(\text{pw}(G) + 1) \rfloor$.
- We prove new bounds for $\chi_{ON}(G)$ with respect to the structural parameters neighborhood diversity and distance to cluster, improving the existing results of Gargano and Rescigno [Theor. Comput. Sci. 2015] and Reddy [Theor. Comput. Sci. 2018], respectively. Furthermore, our techniques also yield improved bounds for the closed neighborhood variant of the problem.
- We also study the partial coloring variant of the CFON coloring problem, which allows vertices to be left uncolored. Let $\chi_{ON}^*(G)$ denote the minimum number of colors required to color G as per this variant. Abel et. al. [SIDMA 2018] showed that $\chi_{ON}^*(G) \leq 8$ when G is planar. They asked if fewer colors would suffice for planar graphs. We answer this question by showing that $\chi_{ON}^*(G) \leq 5$ for all planar G . This approach also yields the bound $\chi_{ON}^*(G) \leq 4$ for all outerplanar G .

All our bounds are a result of constructive algorithmic procedures.

1 Introduction

A *proper coloring* of a graph is an assignment of a color to every vertex of the graph such that adjacent vertices receive distinct colors. Conflict-free coloring

is a variant of the graph coloring problem. A conflict-free coloring of a graph G is a coloring such that for every vertex in G , there exists a uniquely colored vertex in its neighborhood. This problem was first introduced in 2002 by Even, Lotker, Ron and Smorodinsky [1]. This problem was originally motivated by wireless communication systems, where the base stations and clients have to communicate with each other. Each base station is assigned a frequency and if two base stations with the same frequency communicate with the same client, it leads to interference. So for each client, it is ideal to have a base station with a unique frequency. Since each frequency band is expensive, there is a need to minimize the number of frequencies used by the base stations.

Over the past two decades, this problem has been very well studied, see for instance the survey by Smorodinsky [2]. The conflict-free coloring problem has been studied with respect to the open neighborhood and the closed neighborhood. In this paper, we focus on the open neighborhood variant of the problem.

Definition 1 (Conflict-Free Coloring). A CFON coloring of a graph $G = (V, E)$ using k colors is an assignment $C : V(G) \rightarrow \{1, 2, \dots, k\}$ such that for every $v \in V(G)$, there exists an $i \in \{1, 2, \dots, k\}$ such that $|N(v) \cap C^{-1}(i)| = 1$. The smallest number of colors required for a CFON coloring of G is called the CFON chromatic number of G , denoted by $\chi_{ON}(G)$.

The closed neighborhood variant of the problem, CFCN coloring, is obtained by replacing the open neighborhood $N(v)$ by the closed neighborhood $N[v]$ in the above. The corresponding chromatic number is denoted by $\chi_{CN}(G)$.

The CFON coloring problem and many of its variants are known to be NP-complete [3, 4]. It was further shown in [4] that the CFON coloring problem is hard to approximate within a factor of $n^{1/2-\epsilon}$, unless $P = NP$. Since the problem is NP-hard, the parameterized aspects of the problem have been studied. The problems are fixed parameter tractable when parameterized by vertex cover number, neighborhood diversity [4], distance to cluster [5], and more recently, treewidth [6, 7]. This problem has attracted special interest for graphs arising out of intersection of geometric objects, see for instance, [8–10].

The CFON coloring problem is considered as the harder of the open and closed neighborhood variants, see for instance, remarks in [8, 11]. It is easy to construct example graphs G , for which $\chi_{CN}(G) = 2$ and $\chi_{ON}(G) = \Theta(\sqrt{n})$. Pach and Tardos [11] showed that for any graph G on n vertices, the closed neighborhood chromatic number $\chi_{CN}(G) = O(\log^2 n)$. The corresponding best bound [11, 12] for open neighborhood is $\chi_{ON}(G) = O(\sqrt{n})$.

Another variant that has been studied [3] is the partial coloring variant:

Definition 2 (Partial Conflict-Free Coloring). A partial conflict-free coloring on open neighborhood, denoted by $CFON^*$, of a graph $G = (V, E)$ using k colors is an assignment $C : V(G) \rightarrow \{1, 2, \dots, k, \text{unassigned}\}$ such that for every $v \in V(G)$, there exists an $i \in \{1, 2, \dots, k\}$ such that $|N(v) \cap C^{-1}(i)| = 1$.

The corresponding $CFON^*$ chromatic number is denoted $\chi_{ON}^*(G)$.

The key difference between $CFON^*$ coloring and CFON coloring is that in the partial variant, we allow some vertices to be not assigned a color. If a graph can

be CFON* colored using k colors, then all the uncolored vertices can be assigned the color $k + 1$, and thus is a CFON coloring using $k + 1$ colors.

1.1 Our Results and Discussion

In this paper, we obtain improved bounds for $\chi_{ON}(G)$ under different settings. More importantly, all our bounds are a result of constructive algorithmic procedures and hence can easily be converted into respective algorithms. We summarize our results below:

1. In Section 3, we show that $\chi_{ON}(G) \leq \lfloor \frac{5}{3}(\text{pw}(G) + 1) \rfloor$ where $\text{pw}(G)$ denotes the pathwidth of G . The previously best known bound in terms of $\text{pw}(G)$ was $\chi_{ON}(G) \leq 2\text{pw}(G) + 1$, implied by the results in [6].

To the best of our knowledge, this is the first upper bound for $\chi_{ON}(G)$ in terms of pathwidth, which does not follow from treewidth. Our bound follows from an algorithmic procedure and uses an intricate analysis. We are unable to generalize our bound in terms of treewidth because we crucially use a fact (stated as Theorem 6) that applies to path decomposition, but does not seem to apply to tree decomposition. It will be of interest to see if this hurdle can be overcome to obtain an equivalent bound in terms of treewidth.

There are graphs G for which $\chi_{ON}(G) = \text{tw}(G) + 1 = \text{pw}(G)$. It would be interesting to close the gaps between the respective upper and lower bounds.

2. In Section 4, we show that $\chi_{ON}(G) \leq \text{fvs}(G) + 2$, where $\text{fvs}(G)$ denotes the size of a minimum feedback vertex set of G . This bound is tight and is an improvement over the bound $\chi_{ON}(G) \leq \text{fvs}(G) + 3$ by Bodlaender, Kolay and Pieterse [6].
3. In Section 5.1, we give improved bounds with respect to neighborhood diversity parameter. Gargano and Rescigno [4] showed that $\chi_{ON}(G) \leq \chi_{ON}(H) + \text{cl}(G) + 1$ and $\chi_{CN}(G) \leq \chi_{CN}(H) + \text{ind}(G) + 1$. Here H is the type graph of G , while $\text{cl}(G)$ and $\text{ind}(G)$ denote the number of cliques and independent sets respectively in the type partition of G . We present the improvements $\chi_{ON}(G) \leq \chi_{ON}(H) + \text{cl}(G)/2 + 2$ and $\chi_{CN}(G) \leq \chi_{CN}(H) + \text{ind}(G)/3 + 3$.
4. In Section 5.2, we show that $\chi_{ON}(G) \leq \text{dc}(G) + 3$, where $\text{dc}(G)$ is the distance to cluster parameter of G . This is an improvement over the previous bound [5] of $2\text{dc}(G) + 3$. Our bound is nearly tight since there are graphs for which $\chi_{ON}(G) = \text{dc}(G)$. Using a similar approach, we obtain the improved bound $\chi_{CN}(G) \leq \max\{3, \text{dc}(G) + 1\}$.

For the results in terms of parameters neighborhood diversity and distance to cluster, the obvious open questions are to improve the bounds and/or to provide tight examples.

5. When G is planar, we show that $\chi_{ON}^*(G) \leq 5$. This improves the previous best known bound by Abel et al. [3] of $\chi_{ON}^*(G) \leq 8$. The same approach helps us show that $\chi_{ON}^*(G) \leq 4$, when G is an outerplanar graph. These two results are discussed in Section 6.

There are planar graphs G for which $\chi_{ON}^*(G) = 4$, which shows that our bound is nearly tight and leaves a gap of 1 between the upper and lower bounds. It will be of interest to close this gap.

6. For outerplanar graphs G , the bound $\chi_{ON}^*(G) \leq 4$ implies a bound of $\chi_{ON}(G) \leq 5$. We show a better bound of $\chi_{ON}(G) \leq 4$.

2 Preliminaries

In this paper, we consider only simple, finite, undirected and connected graphs. If the graph is not connected, we color each of the components independently. Also, we assume that the graphs do not have isolated vertices as they cannot be CFON colored. The graph induced by a set of vertices V' in G is denoted $G[V']$. For any two vertices $u, v \in V(G)$, the shortest distance between them is denoted $dist(u, v)$. The open neighborhood of v , denoted $N(v)$, is the set of vertices adjacent to v . The closed neighborhood of v , denoted $N[v]$, is $N[v] = N(v) \cup \{v\}$. The degree of a vertex v in the graph is denoted $deg(v)$. The distance, degree and neighborhood restricted to a subgraph H is denoted $dist_H(u, v)$, $deg_H(v)$ and $N_H(v)$ respectively.

We denote the set $\{1, 2, \dots, q\}$ by $[q]$. Throughout this paper, we use the coloring functions $C : V \rightarrow [q]$ and $U : V \rightarrow [q]$ to denote the color assigned to a vertex and a unique color in its neighborhood, respectively. For a vertex $v \in V(G)$, if there exists a vertex $w \in N(v)$ such that $\{x \in N(v) \setminus \{w\} : C(x) = C(w)\} = \emptyset$, then w is called a uniquely colored neighbor of v .

For theorems marked \star , we provide the full proofs in the appendix due to space constraints.

3 Pathwidth

Theorem 3 (Main Pathwidth Result). *Let G be a graph and let $pw(G)$ denote the pathwidth of G . Then there exists a CFON coloring of G using at most $\lfloor \frac{5}{3}(pw(G) + 1) \rfloor$ colors.*

The proof of this theorem will be a constructive procedure that assigns colors to the vertices of G from a set of size $5(pw(G) + 1)/3$. We first formally define pathwidth.

Definition 4 (Path decomposition [13]). *A path decomposition of a graph G is a sequence $\mathcal{P} = (X_1, X_2, \dots, X_s)$ of bags such that, for every $p \in \{1, 2, \dots, s\}$, we have $X_p \subseteq V(G)$ and the following hold:*

- For each vertex $v \in V(G)$, there is a $p \in \{1, 2, \dots, s\}$ such that $v \in X_p$.
- For each edge $\{u, v\} \in E(G)$, there is a $p \in \{1, 2, \dots, s\}$ such that $u, v \in X_p$.
- If $v \in X_{p_1}$ and $v \in X_{p_2}$ for some $p_1 \leq p_2$, then $v \in X_p$ for all $p_1 \leq p \leq p_2$.

The *width* of a path decomposition (X_1, X_2, \dots, X_s) is $\max_{1 \leq p \leq s} \{|X_p| - 1\}$. The *pathwidth* of a graph G , denoted $pw(G)$, is the minimum width over all path decompositions of G . For the purposes of our algorithm, we need the path decomposition to satisfy certain additional properties too.

Definition 5 (Semi-Nice Path Decomposition). A path decomposition $\mathcal{P} = (X_1, X_2, \dots, X_s)$ is called a semi-nice path decomposition if $X_1 = X_s = \emptyset$ and for all $p \in \{2, \dots, s\}$, exactly one of the following hold:

- SN1.** There is a vertex v such that $v \notin X_{p-1}$ and $X_p = X_{p-1} \cup \{v\}$. In this case, we say that X_p introduces v . Further, when X_p introduces v , $N(v) \cap X_p \neq \emptyset$.
- SN2.** There is a vertex v such that $v \in X_{p-1}$ and $X_p = X_{p-1} \setminus \{v\}$. In this case, we say X_p forgets v .
- SN3.** There is a pair of vertices v, \hat{v} such that $v, \hat{v} \notin X_{p-1}$ and $X_p = X_{p-1} \cup \{v, \hat{v}\}$. We call such a bag X_p a special bag that introduces v and \hat{v} . Further, in a special bag X_p that introduces v and \hat{v} , it must be true that $N(v) \cap X_p = \{\hat{v}\}$ and $N(\hat{v}) \cap X_p = \{v\}$.

We first note that the every graph without isolated vertices has a semi-nice path decomposition of width $\text{pw}(G)$.

Theorem 6. Let G be a graph that has no isolated vertices. Then it has a semi-nice path decomposition of width $\text{pw}(G)$.

The proof of the above theorem is deferred to Section 3.1, after the proof of the main theorem of this section.

Algorithm We start with a semi-nice path decomposition $\mathcal{P} = (X_1, X_2, \dots, X_q)$ of width $\text{pw}(G)$. We process each bag in the order X_1, X_2, \dots, X_q . As we encounter each bag, we assign to the vertices in the bag a color $C : V(G) \rightarrow [5(\text{pw}(G) + 1)/3]$. We will also identify a unique color (from its neighborhood) for each vertex $U : V(G) \rightarrow [5(\text{pw}(G) + 1)/3]$. We color the bags such that the below are satisfied:

- Invariant 1.** For any bag X , if $v, v' \in X$, then $C(v) \neq C(v')$.
- Invariant 2.** Suppose we have processed bags X_1 to X_p , where $p \geq 2$. At this point, the induced graph $G[\cup_{1 \leq j \leq p} X_j]$ is CFON colored.
- Invariant 3.** For every vertex v that appears in the bags processed, $U(v)$ is set as $C(w)$ for a neighbor w of v . Once $U(v)$ is assigned, it is ensured that for all “future” neighbors v' of v , $C(v') \neq U(v)$, thereby ensuring that $U(v)$ is retained as a unique color in $N(v)$.

Definitions required for the algorithm: For each bag X , we define the set of *free colors*, as $F(X) = \{U(x) : x \in X\} \setminus \{C(x) : x \in X\}$. That is, $F(X)$ is the set of colors that appear in X as unique colors of vertices in X , but not as colors of any vertex. Further, we partition $F(X)$ into two sets $F_1(X)$ and $F_{>1}(X)$. They are defined as $F_1(X) = \{c \in F(X) : |\{x \in X : U(x) = c\}| = 1\}$ and $F_{>1}(X) = \{c \in F(X) : |\{x \in X : U(x) = c\}| > 1\}$. A vertex v that appears in a bag X is called a *needy vertex* (or simply *needy*) in X , if $U(v) \in F(X)$. For a bag X , we say that a set $S \subseteq X$ is an *expensive subset* if $|\cup_{w \in S} \{C(w), U(w)\}| = 2|S|$.

When going through the sequence of bags in the semi-nice path decomposition, the bags X that forget a vertex only contain vertices that have already been assigned colors and hence no action needs to be taken. When we move from

a bag X' to the next bag X that introduces either one vertex or two vertices, we need to handle the introduced vertices. Let us first consider the bags that introduce one vertex, say v . For all bags that introduce one vertex, we assign $C(v)$ and $U(v)$ as per the below rules.

For bags that introduce one vertex v

Rule 1 for assignment of $C(v)$:

- If there exists a color $c \in F_1(X') \setminus \{U(x) : x \in N(v) \cap X'\}$, then we assign $C(v) = c$. If there are more than one such color c , choose a c such that $|\{x : x \in X', C(U^{-1}(c)) = U(x)\}|$ is minimized. Note that for all $c \in F_1(X')$, there is a unique vertex $w \in X'$ such that $U(w) = c$, and hence $U^{-1}(c)$ is well defined.
- If $F_1(X') \setminus \{U(x) : x \in N(v) \cap X'\} = \emptyset$, we check if there exists a color $c \in F_{>1}(X') \setminus \{U(x) : x \in N(v) \cap X'\}$. If so, we assign $C(v) = c$. If there are multiple such c , then we choose one arbitrarily.
- If $F_1(X') \cup F_{>1}(X') \setminus \{U(x) : x \in N(v) \cap X'\} = \emptyset$, then there are no free colors that can be assigned as $C(v)$. We assign $C(v)$ to be a new color (a color not in $\cup_{x \in X'} \{C(x), U(x)\}$).

Rule 2 for assignment of $U(v)$: We assign $U(v) = C(y)$, where $y \in X'$ is a neighbor of v . Such a y exists by Theorem 6. If v has multiple neighbors, we follow the below priority order:

- If v has needy vertices in X' as neighbors, we choose y as a needy neighbor such that $|\{x : x \in X', U(y) = U(x)\}|$ is minimized.
- If v does not have needy vertices in X' as neighbors, then we choose $y \in X'$ arbitrarily from the set of neighbors of v .

Now let us consider the case where the bag X is a special bag that introduces two vertices v and \hat{v} . We assign $C(v), C(\hat{v}), U(v), U(\hat{v})$ as per the following:

For special bags that introduce two vertices v and \hat{v}

For assignment of $C(v)$ and $C(\hat{v})$: We select one of v and \hat{v} arbitrarily, say v , to be colored first. We use Rule 1 to assign $C(v)$ and then $C(\hat{v})$, in that order. One point to note is that during the application of Rule 1 here, the part $\{U(x) : x \in N(v) \cap X'\}$ will not feature as neither v nor \hat{v} have neighbors in X' .

For assignment of $U(v)$ and $U(\hat{v})$: Assign $U(v) = C(\hat{v})$ and $U(\hat{v}) = C(v)$.

It can easily be checked that the above rules maintain the invariants 1, 2 and 3 stated earlier and hence the algorithm results in a CFON coloring of G . What remains is to show that $5(\text{pw}(G) + 1)/3$ colors are sufficient. We first prove a technical result.

Theorem 7 (Technical Pathwidth Result). *During the course of the algorithm, let k be the size of the largest expensive subset out of all the bags in the path decomposition. Then there must exist a bag of size at least $3k/2$.*

Proof. In the sequence of bags seen by the algorithm, let X be the first bag that has an expensive subset of size k . We show that $|X| \geq 3k/2$. Let $S = \{v_1, v_2, \dots, v_k\} \subseteq X$ be an expensive subset of size k . For each v_i , let $C(v_i) = 2i - 1$ and $U(v_i) = 2i$.

Let X' be the bag that precedes X in the sequence. By the choice of X , no k -expensive subset is present in X' . It follows that $S \not\subseteq X'$. Hence the bag X must introduce a vertex¹ that belongs to S . Without loss of generality, let v_k be this vertex introduced in X . Further wlog, let v_1, \dots, v_r be the needy vertices (in X') of S for some $1 \leq r \leq k$. If none of the vertices in S are needy, then we have that $|X| \geq 2|S| = 2k$ and the theorem holds. So we can assume that $r \geq 1$.

Since the vertices v_1, \dots, v_r are needy in X' , we have $\{2, 4, \dots, 2r\} \subseteq F(X')$. The vertices v_{r+1}, \dots, v_k are not needy because there exist distinct vertices $Z = \{z_{r+1}, \dots, z_k\}$ in the bag X such that $C(z_i) = U(v_i) = 2i$ for $r+1 \leq i \leq k$. We have three cases. In Cases 1 and 2, X is a bag that introduces one vertex v_k . Case 1 is when none of the colors in $F(X')$ was eligible to be assigned as $C(v_k)$. Hence $C(v_k)$ is assigned from outside the set $\cup_{x \in X'} \{C(x), U(x)\}$. Case 2 is when there are eligible colors in $F(X')$, and $C(v_k)$ is chosen from $F(X') \subseteq \cup_{x \in X'} \{C(x), U(x)\}$. Case 3 is when X is a special bag that introduces two vertices.

Case 1: X is a bag that introduces one vertex v_k and $2k - 1 \notin \{U(x) : x \in X'\}$. There is no vertex $x \in X$ with $U(x) = 2k - 1$. To assign a color to v_k , the algorithm chose a new color. This means that $F(X') \setminus \{U(x) : x \in N(v_k) \cap X'\} = \emptyset$. In particular, for each $1 \leq i \leq r$, there² exists $v'_i \in N(v_k) \cap X'$ such that $U(v'_i) = 2i$. Hence the colors $2i$, for $1 \leq i \leq r$ cannot be assigned as $C(v_k)$. By Rule 2, we must set $U(v_k)$ to be the $C(y)$ where y is a needy neighbor of v_k . Hence $C(y) = 2k$.

If $U(y) \notin \{1, 2, 3, \dots, 2k - 2\}$, then $(S \cup \{y\}) \setminus \{v_k\}$ is a k -expensive subset in X' , the predecessor of X . This contradicts the choice of X . Hence we can assume that $U(y) \in \{1, 2, 3, \dots, 2k - 2\}$. Since y is needy in X' , $U(y)$ is not $C(v)$ for any $v \in X'$ and hence we conclude $U(y) \notin \{1, 3, \dots, 2k - 3\}$. Further, colors from $\{2(r+1), 2(r+2), \dots, 2k - 2\}$ appear as $C(z)$ for the vertices $z \in Z$. Hence $U(y) \in \{2, 4, 6, \dots, 2r\}$.

Let $U(y) = 2j$ for some $1 \leq j \leq r$. Notice that $U(v_j) = 2j$ as well, giving us $|\{x : x \in X', U(y) = U(x)\}| \geq 2$.

By Rule 2, we chose $U(v_k) = C(y)$, where y is the needy neighbor that minimizes $|\{x : x \in X', U(y) = U(x)\}|$. We chose y over other needy neighbors v'_1, \dots, v'_r of v_k . Hence there exist r distinct vertices $Y = \{y_1, \dots, y_r\}$ in the bag X , disjoint from S , such that $U(y_i) = U(v'_i) = 2i$ for each $1 \leq i \leq r$.

¹ In the case where X is a special bag that introduces two vertices, at most one of the two introduced vertices can be part of an expensive subset.

² The vertex v'_i may or may not be the same as v_i .

Note that the set $Y \cup Z$ must be disjoint from S , but Y and Z may intersect with each other. Since $|Y| + |Z| = k$, we have $|Y \cup Z| \geq k/2$ and therefore $|X| \geq |S| + |Y \cup Z| \geq 3k/2$.

Case 2 (Proof Sketch): X is a bag that introduces one vertex v_k and $2k - 1 \in \{U(x) : x \in X'\}$. For the sake of brevity and clarity, the full proof of Case 2 is deferred to Appendix A.1. The arguments are similar to the ones used in Case 1, but Case 2 requires a lengthier treatment. We give a sketch of the proof here.

In this case, $C(v_k)$ is chosen from $F(X')$. That is, $C(v_k)$ is chosen as $U(w) = 2k - 1$, for a vertex w that is needy in X' , chosen according to Rule 1. It can be first established that $C(w) \in \{2(r + 1), 2(r + 2), \dots, 2k - 2\}$.

We know that $\{2, 4, \dots, 2r\} \subseteq F(X')$. Let $\ell = |\{2, 4, \dots, 2r\} \cap \{U(x) : x \in N(v_k) \cap X'\}|$. The color $2k - 1$ was chosen as $C(v_k)$ over the other colors in $F(X')$. In particular, it was chosen over the $r - \ell$ colors in $\{2, 4, \dots, 2r\} \setminus \{U(x) : x \in N(v_k) \cap X'\}$. This is used to show the existence of a set $W \subseteq X$ of size $r - \ell$ that is disjoint from S .

Now we study why $U(v_k)$ was assigned as $2k$. As per Rule 2, $U(v_k)$ was assigned as $C(y)$, where y is a needy neighbor of v_k . There exists at least ℓ needy neighbors of v_k . This is used to establish the existence of a set $Y \subseteq X$ of size ℓ that is disjoint from S and W .

We thus have $|W \cup Y| = r$ and $|Z| = k - r$. The sets $W \cup Y$ and Z are both disjoint from S , but need not be disjoint from each other. Hence $|X| \geq |S| + |W \cup Y \cup Z| \geq 3k/2$.

Case 3 (Proof Sketch): X is a special bag that introduces v_k and \hat{v}_k . As in Case 2, the full proof of Case 3 is given in Appendix A.2. We give a sketch of the proof here.

If $F(X') = \emptyset$, then none of the $k - 1$ vertices in $S \cap X'$ are needy in X' . Hence $|X'| \geq 2(k - 1)$. This implies that $|X| \geq 2(k - 1) + 2 = 2k$ and we are done.

Else, $|F(X')| \geq 1$. Let us first note that since S is an expensive subset, so is $S \cup \{\hat{v}_k\} \setminus \{v_k\}$. Since $|F(X')| \geq 1$, at least one of $C(v_k)$ or $C(\hat{v}_k)$ will be chosen from $F(X')$. Without loss of generality, let v_k be a vertex such that $C(v_k) \in F(X')$. Let $C(v_k) = U(w) = 2k - 1$, where w is a needy vertex in X' , chosen according to Rule 1. The rest of the arguments are very similar to the arguments in the proof of Case 2.

We finally establish the existence of a set $W \subseteq X$ such that W is disjoint from S and $|W| = r$. Hence $|X| \geq |S| + |W \cup Z| \geq 3k/2$. \square

Now we prove the main theorem of this section.

Proof (Proof of Theorem 3). We apply the algorithm in a nice path decomposition of G , which satisfies the condition in Theorem 6. As stated before, the correctness follows by the stated invariants, and what remains to be shown is the bound on the number of colors necessary.

Consider any bag X of the path decomposition. Then we have

$$|\cup_{w \in X} \{C(w), U(w)\}| = |X| + |\text{Extra}(X)|,$$

where $\text{Extra}(X)$ denotes the set of colors that feature as unique colors, but not as colors of vertices in X .

We construct a subset Y of X as follows. For each color in $\text{Extra}(X)$, we include exactly one vertex y in Y such that $U(y)$ is that color. We have $|Y| = |\text{Extra}(X)|$. We also have that Y is an expensive subset of X . Since no bag is of size bigger than $\text{pw}(G) + 1$, it follows that by Theorem 7 that $|Y| \leq 2(\text{pw}(G) + 1)/3$. Since $|Y|$ is an integer, we can say $|Y| \leq \lfloor 2(\text{pw}(G) + 1)/3 \rfloor$. Hence

$$|\cup_{w \in X} \{C(w), U(w)\}| \leq |X| + \lfloor 2(\text{pw}(G) + 1)/3 \rfloor.$$

In the algorithm, we need to add a new color to the bag only when a bag X is followed by another bag that introduces a vertex. Hence we may require one additional color, which brings the maximum number of colors needed to $|X| + \lfloor 2(\text{pw}(G) + 1)/3 \rfloor + 1 \leq \text{pw}(G) + \lfloor 2(\text{pw}(G) + 1)/3 \rfloor + 1 = \lfloor 5(\text{pw}(G) + 1)/3 \rfloor$. \square

3.1 Proof of Theorem 6

A path decomposition (X_1, X_2, \dots, X_s) is called a *nice path decomposition* if the following hold:

- $X_1 = X_s = \emptyset$.
- For all $p \in \{2, 3, \dots, s\}$, there is a vertex v such that either $v \notin X_{p-1}$ and $X_p = X_{p-1} \cup \{v\}$, or $v \in X_{p-1}$ and $X_p = X_{p-1} \setminus \{v\}$. In the former case, we say X_p *introduces* v , and in the latter case we say X_p *forgets* v .

It is known [13] that every graph G has a nice path decomposition of width equal to $\text{pw}(G)$, and that every nice path decomposition has exactly $2|V(G)| + 1$ bags.

Consider a nice path decomposition of the graph G . If all the vertices have a neighbor in the bag that introduces them, then the nice path decomposition is itself a semi-nice path decomposition, and we are done. Otherwise, we explain how to convert the given nice path decomposition into a semi-nice path decomposition. We say that a bag X is a *violating bag* if it introduces a vertex v and $X \cap N(v) = \emptyset$. The violating bags do not follow the rules SN1, SN2, or SN3 from Definition 5. Instead they follow the below rule SN1'.

SN1'. There is a vertex v such that $v \notin X_{p-1}$ and $X_p = X_{p-1} \cup \{v\}$. In this case, we say that X_p *introduces* v . Further, when X_p introduces v , $N(v) \cap X_p = \emptyset$.

We say that a path decomposition is a *t-violating semi-nice path decomposition* if there are t violating bags and the rest of the bags obey one of the rules SN1, SN2 or SN3 from Definition 5.

We “fix” each violating bag by modifying the path decomposition. The fix involves delaying the introduction of a vertex until it has a neighbor, and possibly creating a “special bag” that introduces two vertices. Throughout the fix-up process, the path decomposition in hand will be a *t-violating semi-nice decomposition*, with every step of the fix-up decrementing t by one. We now explain the fix-up process.

Fix-up Process: Given a t -violating semi-nice path decomposition $\mathcal{P} = (X_1, X_2, \dots, X_s)$, we explain how to obtain a $(t-1)$ -violating semi-nice path decomposition \mathcal{P}' .

Let X_{p_1} be a violating bag that introduces the vertex v , which is forgotten by the bag X_{p_2} . By assumption, $X_{p_1} \cap N(v) = \emptyset$. Let X_q be the first bag in the sequence that contains a neighbor of v . Since G does not have isolated vertices, $N(v)$ is non-empty, and hence $p_1 < q < p_2$. Let $\hat{v} \in X_q$ be a neighbor of v . We have two cases.

Case 1: $|N(\hat{v}) \cap X_q| > 1$. That is, \hat{v} has other neighbors in X_q apart from v . We consider the following modified sequence \mathcal{P}' :

$$X_1, \dots, X_{p_1-1}, X_{p_1+1} \setminus \{v\}, X_{p_1+2} \setminus \{v\}, \dots, X_{q-1} \setminus \{v\}, X_q \setminus \{v\}, X_q, X_{q+1}, \dots, X_s$$

That is, we delay the introduction of v till its first neighbor, \hat{v} , has been introduced. It can be verified that the above sequence \mathcal{P}' is a path decomposition of the same graph G with width no more than the width of \mathcal{P} . In the new sequence \mathcal{P}' , v is introduced by X_q which is not a violating bag in \mathcal{P}' . Below, we explain that the fix-up process has not introduced any new violations. Since v sees \hat{v} in X_q for the first time, it follows that the bag X_q introduces \hat{v} in \mathcal{P} .

We first note that X_q was not a special bag in \mathcal{P} . To see why, let us assume the contrary. Let X_q be a special bag in \mathcal{P} that introduces the vertices \hat{v} and w . If so, $N(\hat{v}) \cap X_q = \{w\}$ as per SN3. Hence $v \notin X_q$. So we can conclude that X_q is not a special bag, and therefore X_q introduces just one vertex \hat{v} as per SN1. This means that v cannot have any other neighbors in X_q apart from \hat{v} . Hence \hat{v} is the only vertex that loses a neighbor from its introducing bag due to the fix-up process. However, since \hat{v} has other neighbors in X_q apart from v , it does not result in a violation in \mathcal{P}' .

Case 2: $|N(\hat{v}) \cap X_q| = 1$. That is, v is the lone neighbor of \hat{v} in X_q . Since X_q is the first bag in \mathcal{P} that contains a neighbor of v , it follows that X_q introduces \hat{v} . Since v is the lone neighbor of \hat{v} in X_q , it follows that X_q is not a special bag in \mathcal{P} . Hence \hat{v} is the lone neighbor of v in X_q . We consider the following sequence \mathcal{P}' :

$$X_1, \dots, X_{p_1-1}, X_{p_1+1} \setminus \{v\}, X_{p_1+2} \setminus \{v\}, \dots, X_{q-1} \setminus \{v\}, X_q, X_{q+1}, \dots, X_s$$

We introduce v together with \hat{v} in the bag X_q . We have already seen that $N(v) \cap X_q = \{\hat{v}\}$ and $N(\hat{v}) \cap X_q = \{v\}$. Thus X_q becomes a special bag in \mathcal{P}' that introduces v and \hat{v} . No other violations have been introduced by this because v does not have any neighbors in $X_{p_1+1}, X_{p_1+1}, \dots, X_{q-1}$.

Thus by repeating this fix-up process for each of the violations, we can convert the given nice path decomposition into a semi-nice path decomposition. \square

4 Feedback Vertex Set

Definition 8 (Feedback Vertex Set). Let $G = (V, E)$ be an undirected graph. A feedback vertex set (FVS) is a set of vertices $S \subseteq V$, removal of which from the graph G makes the remaining graph $(G[V \setminus S])$ acyclic. The size of a smallest such set S is denoted as $\text{fvs}(G)$.

Theorem 9. $\chi_{ON}(G) \leq \text{fvs}(G) + 2$.

The following graph (as observed in [6]), shows that the above theorem is tight. Let K_n^* be the graph obtained by starting with the clique on n vertices, and subdividing each edge with a vertex. Then K_n^* has an FVS of size $n - 2$, and it can be seen that $\chi_{ON}(K_n^*) = n$.

The proof of this theorem is through a constructive process to CFON color the vertices of the graph G , given a feedback vertex set F of G . By definition, $G[V \setminus F]$ is a collection of trees.

Each tree T in $G[V \setminus F]$ is rooted at an arbitrary vertex r_T . If $|V(T)| \geq 2$, we choose a neighbor of r_T and call it the *special vertex* in T , denoted by s_T . Let v be a vertex not in T . The *deepest neighbor* of v in T , denoted by $\text{deep}_T(v)$, is a vertex $w \in V(T) \cap N(v)$ such that $\text{dist}_T(r_T, w)$ is maximized. If there are multiple such vertices at the same distance, the deepest neighbor is chosen to be a vertex which is not the special vertex s_T .

Lemma 10. *Let T be a tree with $|V(T)| \geq 2$. Then $\chi_{ON}(T) \leq 2$.*

Proof. We assign colors $C : V(T) \rightarrow \{1, 2\}$ in the following manner.

- Assign $C(r_T) = 1$ and $C(s_T) = 2$.
- For each vertex $v \in N_T(r_T) \setminus \{s_T\}$, assign $C(v) = 1$.
- For the remaining vertices $v \in V(T)$, assign $C(v) = \{1, 2\} \setminus C(w)$, where w is the grandparent of v .

For each vertex $v \in V(T) \setminus \{r_T\}$, the uniquely colored neighbor is its parent. For r_T , the uniquely colored neighbor is s_T . This is a CFON 2-coloring of T . \square

We first prove a special case of Theorem 9.

Lemma 11. *Let $G = (V, E)$ be a graph and $F \subseteq V$ be a feedback vertex set with $|F| = 1$. Then G can be CFON colored using 3 colors.*

Proof. Let $F = \{v\}$. First using Lemma 10, we color all the trees $T \subseteq G[V \setminus F]$ using the colors 2 and 3, whenever $|V(T)| \geq 2$. All the singleton components of $G[V \setminus F]$ are assigned the color 2. We assign $C(v) = 1$. Now all the vertices, except possibly v , have a uniquely colored neighbor. We explain how to fix this and obtain a CFON coloring.

- **Case 1: There exists a singleton component** $\{w\} \subseteq G[V \setminus F]$.
Reassign $C(w) = 1$.
- **Case 2: Else, if there exists a component** $T \subseteq G[V \setminus F]$, **such that either (i) $\text{deep}_T(v) \neq s_T$ or (ii) $\text{deep}_T(v) = s_T$ and $\{r_T, v\} \notin E(G)$.**
Reassign $C(\text{deep}_T(v)) = 1$.
- **Case 3: Else, for each component** $T \subseteq G[V \setminus F]$, $N(v) \cap V(T) = \{r_T, s_T\}$.
If there exists a component $T \subseteq G[V \setminus F]$, such that $|V(T)| \geq 3$, choose a vertex $w \in V(T) \setminus \{r_T, s_T\}$ and set w as the new root of T . Reassign s_T and the colors of $V(T)$ accordingly. Doing so will ensure that $\text{deep}_T(v) \neq s_T$. We apply Case 2.
Else, for all the components $T \subseteq G[V \setminus F]$, we have $|V(T)| = 2$. Choose a component $T' \subseteq G[V \setminus F]$. For all the other vertices $w \in V \setminus (\{v\} \cup V(T'))$, reassign $C(w) = 2$.

All the trees in $G[V \setminus F]$ are CFON colored as per the earlier described procedure. Even after reassigning some colors, they remain CFON colored. The vertex v sees another vertex w , with $C(w) = 1$ if in Case 1 or 2. In the last case, v sees a unique vertex that is colored 3. \square

4.1 Proof of Theorem 9

When $|F| = 1$, three colors are sufficient to CFON color G by Lemma 11. Now, we consider the case when $|F| \geq 2$. We assign colors $C : V(G) \rightarrow [|F| + 2]$ in such a way that G is CFON colored. First by Lemma 10, we color all the components $T \subseteq G[V \setminus F]$ with $|V(T)| \geq 2$, using the colors $|F| + 1$ and $|F| + 2$.

During the algorithm, we will keep track of a color which when assigned to the isolated vertices in the feedback vertex set does not change the unique color in the neighborhood of the already colored vertices. We call this color a *free color* and denote this by c' , initialized to 0.

Let $F = \{v_1, v_2, \dots, v_{|F|}\}$. Let $Y = \{v_i \in F : \deg_F(v_i) \geq 1\}$. For each $v_i \in Y$, assign $C(v_i) = i$. Note that for every $v_i \in Y$, there is at least one uniquely colored neighbor in Y . If $Y \neq \emptyset$, choose an arbitrary vertex $v_i \in Y$ and set $c' = i$.

Now the vertices of Y and components $T \subseteq G[V \setminus F]$ with $|V(T)| \geq 2$ are colored and have a uniquely colored neighbor. What remains are the vertices in $F \setminus Y$ and singleton components of $G[V \setminus F]$. Below we explain how to color them in phases.

Recall that $G[F \setminus Y]$ is an independent set.

Case 1: Singleton component $\{w\} \subseteq G[V \setminus F]$ where w has at least 1 uncolored neighbor.

- Let $v_{i_1}, v_{i_2}, \dots, v_{i_m} \in F \setminus Y$ be the uncolored neighbors of w , where $m \geq 1$.
- Assign $C(v_{i_1}) = C(w) = i_1$ and $C(v_{i_j}) = i_2$ for all $2 \leq j \leq m$. The uniquely colored neighbor for w is v_{i_1} and for all v_{i_j} , it is the vertex w .
- The free color is set as $c' = i_1$.

Case 2: Singleton component $\{w\} \subseteq G[V \setminus F]$, where all of $N(w)$ is colored and w has no uniquely colored neighbor.

This means that $|N(w)| \geq 2$, and every color in $N(w)$ appears at least twice. Choose two vertices $v_{i_1}, v_{i_2} \in N(w)$ such that $C(v_{i_1}) = C(v_{i_2})$. It must be the case that at least one of the colors i_1, i_2 does not appear in $N(w)$. Without loss of generality, let it be i_1 .

- Reassign $C(v_{i_1}) = i_1$. Assign $C(w) = |F| + 1$. The uniquely colored neighbor for w is v_{i_1} . Notice that all the vertices in $N(w)$ would have received their uniquely colored neighbors when they were assigned a color.
- The free color is set as $c' = i_1$.

Now, all the singleton components $\{w\} \subseteq V \setminus F$ have uniquely colored neighbors, but not all of them may be colored. Assign all the uncolored singleton components the color $|F| + 1$. What remains to be addressed are the remaining uncolored vertices in $F \setminus Y$. These vertices do not have any singleton components of $G[V \setminus F]$ as neighbors. We assign colors if the below Cases 3 or 4 apply.

Case 3: A component $T \subseteq G[V \setminus F]$ such that s_T has at least two uncolored vertices in $F \setminus Y$ as neighbors.

- Let $v_{i_1}, v_{i_2}, \dots, v_{i_m} \in F \setminus Y$ be the uncolored neighbors of s_T , with $m \geq 2$.
- Reassign $C(s_T) = i_1$ and assign $C(v_{i_j}) = i_2$ for all $1 \leq j \leq m$. The vertex s_T serves as the uniquely colored neighbor for the vertices v_{i_j} .
- The free color is set as $c' = i_2$.

Case 4: There exists an uncolored vertex v_i and component $T \subseteq G[V \setminus F]$, such that either (i) $deep_T(v_i) \neq s_T$ or (ii) $deep_T(v_i) = s_T$ and $\{r_T, v_i\} \notin E(G)$.

- Reassign $C(deep_T(v_i)) = i$ and assign $C(v_i) = i$. The vertex $deep_T(v_i)$ serves as the uniquely colored neighbor for v_i .
- The free color is set as $c' = i$.

Case 5 : There exists an uncolored vertex v_i such that for each component $T \subseteq G[V \setminus F]$, either $N(v_i) \cap V(T) = \{r_T, s_T\}$ or $N(v_i) \cap V(T) = \emptyset$. We make use of the free color c' obtained from the previous cases. In this case, we reassign $C(s_T) = i$ and assign $C(v_i) = c'$. The vertex s_T will serve as the uniquely colored neighbor for v_i .

Now we explain why we must have a non-zero free color. If $c' = 0$, we have that $Y = \emptyset$ and none of the previous cases have been applicable. That is:

1. $Y = \emptyset$.
2. There are no singleton components in $G[V \setminus F]$.
3. For each vertex $v_i \in F$ and for each component $T \subseteq G[V \setminus F]$, either $N(v_i) \cap V(T) = \{r_T, s_T\}$ or $N(v_i) \cap V(T) = \emptyset$.
4. For each component $T \subseteq G[V \setminus F]$, $|N(s_T) \cap F| \leq 1$.

Since $|F| \geq 2$, let us consider $v_1, v_2 \in F$. Notice that due to the above, it is not possible to have a path from v_1 to v_2 in G . This means that G is not connected. This is a contradiction. Thus we must have $c' \neq 0$.

We have described a procedure to obtain a CFON coloring that uses $|F| + 2$ colors. By setting F to be a minimum sized FVS, we get $\chi_{ON}(G) \leq \text{fvs}(G) + 2$. \square

5 Neighborhood Diversity & Distance to Cluster

In this section, we give improved bounds for $\chi_{ON}(G)$ and $\chi_{CN}(G)$ with respect to the parameters neighborhood diversity and distance to cluster.

5.1 Neighborhood Diversity

Definition 12 (Neighborhood Diversity [4]). Give a graph $G = (V, E)$, two vertices $v, w \in V$ have the same type if $N(v) \setminus \{w\} = N(w) \setminus \{v\}$. A graph G has neighborhood diversity at most t if $V(G)$ can be partitioned into t sets V_1, V_2, \dots, V_t , such that all the vertices in each V_i , $1 \leq i \leq t$ have the same type. The partition $\{V_1, V_2, \dots, V_t\}$ is called the type partition of G .

It can be inferred from the above definition that all vertices in a V_i either form a clique or an independent set, $1 \leq i \leq t$. For two types V_i, V_j , either each vertex in V_i is neighbor to each vertex in V_j , or no vertex in V_i is neighbor to any vertex in V_j . This leads to the definition of the *type graph* $H = (\{1, 2, \dots, t\}, E_H)$, where $E_H = \{\{i, j\} : 1 \leq i < j \leq t, \text{ each vertex in } V_i \text{ is a neighbor of each vertex in } V_j\}$.

In the above, $cl(G)$ and $ind(G)$ respectively denote the number of V_i 's that form a clique and independent set in the type partition $\{V_1, V_2, \dots, V_t\}$. Gargano and Rescigno [4] showed that the CFON and CFCN variants are fixed parameter tractable with respect to neighborhood diversity. They also obtained the bounds $\chi_{ON}(G) \leq \chi_{ON}(H) + cl(G) + 1$ and $\chi_{CN}(G) \leq \chi_{CN}(H) + ind(G) + 1$. We improve both these bounds.

Theorem 13. $\chi_{ON}(G) \leq \chi_{ON}(H) + \frac{cl(G)}{2} + 2$.

Theorem 14 (\star). $\chi_{CN}(G) \leq \chi_{CN}(H) + \frac{ind(G)}{3} + 3$.

We prove Theorem 13 below. The proof of Theorem 14 uses similar ideas and is presented in Appendix B.

Proof (Proof of Theorem 13). To begin with, we CFON color the type graph H using $\chi_{ON}(H)$ colors. Let $C_H : V_H \rightarrow [\chi_{ON}(H)]$ be that coloring and $U_H : V_H \rightarrow [\chi_{ON}(H)]$ be the corresponding assignment of unique colors. Now, we derive a coloring $C : V(G) \rightarrow \{0, 1, 2, \dots, s, s+1\}$ from C_H , where $s = \chi_{ON}(H) + \frac{cli(G)}{2}$. Also we identify a unique color in the neighborhood of each vertex, denoted $U : V(G) \rightarrow \{0, 1, 2, \dots, s, s+1\}$. Let V_1, V_2, \dots, V_t be the type partition of V . We assign colors to the vertices as follows: For each V_i , choose a representative vertex $r_i \in V_i$ and assign $C(r_i) = C_H(i)$. For each V_i , for all vertices $x \in V_i \setminus \{r_i\}$, we assign $C(x) = 0$. We make the below observations.

- Each of the vertices r_i has a uniquely colored neighbor, as C_H is a CFON coloring of H .
- Let V_i be an independent set, and let r_j be the uniquely colored neighbor of r_i . For each $x \in V_i$, r_j serves as the uniquely colored neighbor.
- If V_i is a clique and $U_H(i) \neq C_H(i)$, the uniquely colored neighbor of r_i serves as the uniquely colored neighbor for all vertices in V_i .

What remains to be handled are the type sets V_i which are cliques and $U_H(i) = C_H(i)$. We call these type sets as *bad sets*. We do not consider the singleton V_i 's as bad sets. All the representative vertices r_i see a uniquely colored neighbor, regardless of whether V_i is bad or not. Note that, once a bad set V is fixed, we no longer call it a bad set.

Let A refer to the following set of colors: $A = \{\chi_{ON}(H) + 1, \chi_{ON}(H) + 2, \dots, \chi_{ON}(H) + cli(G)/2\}$. None of the colors from A have been used till now. **Reduction of bad sets:** If there exists a V_i (not necessarily a bad set) that has at least 2 bad sets as neighbors, we do the following. Let $V_{i_1}, V_{i_2}, \dots, V_{i_m}$ be the bad sets adjacent to V_i . Then we reassign $C(r_i) = c$, where $c \in A$ is a color that has not been used till now. The vertex r_i will serve as the uniquely colored

neighbor for all vertices in $V_{i_1}, V_{i_2}, \dots, V_{i_m}$ as well as the vertices in $V_i \setminus \{r_i\}$. Thus after this operation, none of $V_{i_1}, V_{i_2}, \dots, V_{i_m}$ and V_i are bad sets.

We apply the above reduction operation as much as possible, choosing a new color from A each time. After that, each set V_i is adjacent to at most one bad set. This leaves us with the following two cases.

- **Case 1: Bad sets V_i and V_j which are neighbors, each of which is not neighbors to any other bad sets.**

Reassign $C(r_i) = s+1$. The uniquely colored neighbor of r_i remains the same. And r_i becomes the uniquely colored neighbor for all vertices $x \in V_i \cup V_j \setminus \{r_i\}$. Note that any set V_k that relied on V_i for its unique color, can continue to do so. This is because V_k sees at most one bad set after the repeated application of the reduction operation.

- **Case 2: Bad set V_i , which has no neighboring bad set.**

Let V_j be the neighboring set of V_i such that $C(r_i) = C(r_j)$. We reassign $C(r_i) = s+1$.

Every vertex in V_i has r_j as its uniquely colored neighbor. As in the previous case, any set that relied on V_i for its unique color can continue to do so.

The above is a CFON coloring. We use $\chi_{ON}(H)$ colors to color the representative vertices of each V_i . Each application of the reduction operation needs one new color from A to handle at least two bad sets. Since each bad set is a clique, the number of extra colors needed is at most $cl(G)/2$. Taking the colors $\{0, s+1\}$ into account, the total number of colors used is $\chi_{ON}(H) + cl(G)/2 + 2$. \square

5.2 Distance to Cluster

Definition 15 (Distance to Cluster). Let $G = (V, E)$ be a graph. The distance to cluster of G , denoted $dc(G)$, is the size of the smallest set $X \subseteq V$ such that $G[V \setminus X]$ is a disjoint union of cliques.

Reddy [5], studied the CFCN and the CFON variants with respect to the distance to cluster parameter, $dc(G)$. They showed that $\chi_{ON}(G) \leq 2dc(G) + 3$ and $\chi_{CN}(G) \leq dc(G) + 2$. We give the following improved bounds.

Theorem 16. $\chi_{ON}(G) \leq dc(G) + 3$.

Theorem 17 (*). $\chi_{CN}(G) \leq \max\{3, dc(G) + 1\}$.

For the subdivided clique K_n^* , we have $\chi_{ON}(K_n^*) = dc(K_n^*) = n$. Hence Theorem 16 is nearly tight. We prove Theorem 16 below. The proof of Theorem 17 uses similar ideas and is presented in Appendix C.

Proof (Proof of Theorem 16). Let $dc(G) = d$. That is, there is a set $X \subseteq V(G)$, with $|X| = d$ such that $G[V \setminus X]$ is a disjoint union of cliques.

If $X = \emptyset$, the graph G is a clique because we only consider connected graphs. A clique can be CFON colored using 3 colors. Else, we have $|X| \geq 1$. Let $X = \{v_1, v_2, \dots, v_d\}$. Then $G[V \setminus X] = K_1 \cup K_2 \cdots \cup K_t$ is a disjoint union of cliques.

Below, we explain how to assign colors, $C : V(G) \rightarrow [d+3]$ such that every vertex has a uniquely colored neighbor. We apply the following rules:

1. Let $Y = \{v_i \in X : \deg_X(v_i) \geq 1\}$. For all $v_i \in Y$, assign $C(v_i) = i$.
Now every vertex in Y is colored and has a uniquely colored neighbor.
2. For each of the singleton cliques $K_j = \{w\}$, we do the following.
 - **Case 2(a): The vertex w has at least 1 uncolored neighbor.**
Let $v_{i_1}, v_{i_2}, \dots, v_{i_m} \in X$ be the uncolored neighbors of w , with $m \geq 1$. Assign $C(v_{i_1}) = C(w) = i_1$ and $C(v_{i_\ell}) = d+1$, for all $2 \leq \ell \leq m$. All the vertices in $N(w) \cup \{w\}$ see the color i_1 exactly once in their neighborhood. We will not be assigning the color i_1 for any other vertices henceforth.
 - **Case 2(b): All vertices in $N(w)$ are colored.**
The assignment of colors in the previous case may lead us to this case. If w already sees a uniquely colored neighbor, then we set $C(w) = d+1$. If w has no uniquely colored neighbor, we choose two vertices $v_{i_1}, v_{i_2} \in N(w)$ such that $C(v_{i_1}) = C(v_{i_2})$. Since the only color that is being reused in X is $d+1$, we have $C(v_{i_1}) = C(v_{i_2}) = d+1$. Reassign $C(v_{i_1}) = i_1$. Assign $C(w) = d+1$. Here the color i_1 will be the unique color in the neighborhood of w and this color will not be used in further coloring.
 After this step, all the singleton cliques K_j and their neighboring vertices are colored and also have a uniquely colored neighbor.
3. For each uncolored $v_i \in X \setminus Y$ that does not have a uniquely colored neighbor, we choose a vertex $w \in N(v_i)$ and assign $C(w) = i$. The color i is the unique color in v_i 's neighborhood. And the color i is not used in further coloring.
4. For all the remaining uncolored $v_i \in X \setminus Y$, assign $C(v_i) = d+1$. Recall that v_i is not colored in Step 3 because it has a uniquely colored neighbor.
Now, all the vertices in X are colored and have uniquely colored neighbors. What remains to be colored are the cliques of size at least 2.
5. For each clique K_j with $|K_j| \geq 2$, we note that there may already be some colored vertices in K_j as a result of Step 3. These colors appear exactly once in the graph. We do the following:
 - If K_j has at least 2 colored vertices, color the remaining vertices (if any) with $d+1$.
 - Else, if K_j has exactly 1 colored vertex, choose an uncolored vertex and assign $d+2$. Color the remaining vertices (if any) with $d+1$.
 - Else, choose 2 vertices from K_j and assign the colors $d+2$ and $d+3$. Color the remaining vertices (if any) with $d+1$.

□

6 CFON* Coloring of Planar Graphs

Definition 18 (Planar and Outerplanar graphs). A planar graph is a graph that can be drawn in \mathbb{R}^2 (a plane) such that the edges do not cross each other in the drawing. An outerplanar graph is a planar graph that has a drawing in a plane such that all the vertices of the graph belong to the outer face.

Abel et. al. showed [3] that eight colors are sufficient for CFON* coloring of a planar graph. In this section, we improve the bound to five colors.

We need the following definition:

Definition 19 (Maximal Distance-3 Set). For a graph $G = (V, E)$, a maximal distance-3 set is a set $S \subseteq V(G)$ that satisfies the following:

1. For every pair of vertices $w, w' \in S$, we have $\text{dist}(w, w') \geq 3$.
2. For every vertex $w \in S$, $\exists w' \in S$ such that $\text{dist}(w, w') = 3$.
3. For every vertex $x \notin S$, $\exists x' \in S$ such that $\text{dist}(x, x') < 3$.

The set S is constructed by initializing $S = \{v\}$ where v is an arbitrary vertex. We proceed in iterations. In each iteration, we add a vertex w to S if (1) for every v already in S , $\text{dist}(v, w) \geq 3$, and (2) there exists a vertex $w' \in S$ such that $\text{dist}(w, w') = 3$. We repeat this until no more vertices can be added.

The main component of the proof is the construction of an auxiliary graph G' from the given graph G .

Construction of G' : The first step is to pick a maximal distance-3 set V_0 . Notice that any distance-3 set is an independent set by definition. We let V_1 denote the neighborhood of V_0 . More formally, $V_1 = \{w : \{w, w'\} \in E(G), w' \in V_0\}$. Let V_2 denote the remaining vertices i.e., $V_2 = V \setminus (V_0 \cup V_1)$.

We note the following properties satisfied by the above partitioning of $V(G)$.

1. The set V_0 is an independent set.
2. For every vertex $w \in V_1$, there exists a unique vertex $w' \in V_0$ such that $\{w, w'\} \in E(G)$. This is because if there are two such vertices, this will violate the distance-3 property of V_0 .
3. Every vertex in V_0 has a neighbor in V_1 . If there exists $v \in V_0$ without a neighbor in V_1 , then v is an isolated vertex. By assumption, G does not have isolated vertices.
4. There are no edges from V_0 to V_2 .
5. Every vertex in V_2 has a neighbor in V_1 , and is hence at distance 2 from some vertex in V_0 . This is due to the maximality of the distance-3 set V_0 .

Now we define $A = V_0 \cup V_2$. We first remove all the edges of $G[V_2]$ making A an independent set. For every vertex $v \in A$ we do the following: we identify an arbitrary neighbor $f(v) \in N(v) \subseteq V_1$. Then we contract the edge $\{v, f(v)\}$. That is, we first identify vertex v with $f(v)$. Then for every edge $\{v, v'\}$, we add an edge $\{f(v), v'\}$. The resulting graph is G' .

Theorem 20. If G is a planar graph, $\chi_{ON}^*(G) \leq 5$.

Proof. Let G be a planar graph. We first construct the graph G' as above. Since the steps for constructing G' involve only edge deletion and edge contraction, G' is also a planar graph. By the planar four-color theorem [14], there is an assignment $C : V(G') \rightarrow \{2, 3, 4, 5\}$ such that no two adjacent vertices of G' are assigned the same color. Now we have colored all the vertices in $V(G') = V_1$

Now, we extend C to get a CFON* coloring for G . For all vertices $v \in V_0$, we assign $C(v) = 1$. The vertices in V_2 are not assigned a color.

We will show that C is indeed a CFON* coloring of G . Consider a vertex $v \in A$ which is contracted to a neighbor $f(v) = w \in V_1$. The color assigned to

w is distinct from all w 's neighbors in G' . Hence the color assigned to w is the unique color among the neighbors of v in G .

For each vertex $w \in V_1$, w is a neighbor of exactly one vertex $v \in V_0$. Every vertex $v \in V_0$ is colored 1, which is different from all the colors assigned to the neighbors of w in G' . \square

Outerplanar graphs have a proper coloring using three colors. By argument analogous to Theorem 20, we infer the following.

Corollary 21. *If G is an outerplanar graph, $\chi_{ON}^*(G) \leq 4$.*

For outerplanar graphs, a CFON* coloring using 4 colors implies a CFON coloring using 5 colors. However, we can show the following improved bound. For the sake of clarity, we provide the full proof of the below theorem in Appendix D and give a sketch below.

Theorem 22 (\star). *If G is an outerplanar graph, $\chi_{ON}(G) \leq 4$.*

Proof (Proof Sketch of Theorem 22). Theorem 22 is proved using a two-level induction process. The first level is using a *block decomposition* of the graph. Any connected graph can be viewed as a tree of its constituent blocks. We color the blocks in order so that when we color a block, at most one of its vertices is previously colored. Each block is colored without affecting the color of the already colored vertex. The second level of the induction is required for coloring each of the blocks. We use ear decomposition on each block and color the faces of the block in sequence. However, the proof is quite technical and involves several cases of analysis at each step. \square

Acknowledgments: We would like to thank I. Vinod Reddy for suggesting the problem, Rogers Mathew and N. R. Aravind for helpful discussions and the anonymous reviewer who pointed out an issue with the proof of Theorem 6.

References

1. G. Even, Z. Lotker, D. Ron, S. Smorodinsky, Conflict-free colorings of simple geometric regions with applications to frequency assignment in cellular networks, *SIAM J. Comput.* 33 (1) (2004) 94–136. doi:10.1137/S0097539702431840.
2. S. Smorodinsky, Conflict-Free Coloring and its Applications, Springer Berlin Heidelberg, Berlin, Heidelberg, 2013, pp. 331–389. doi:10.1007/978-3-642-41498-5_12.
3. Z. Abel, V. Alvarez, E. D. Demaine, S. P. Fekete, A. Gour, A. Hesterberg, P. Keldenich, C. Scheffer, Conflict-free coloring of graphs, *SIAM Journal on Discrete Mathematics* 32 (4) (2018) 2675–2702. doi:10.1137/17M1146579.
4. L. Gargano, A. A. Rescigno, Complexity of conflict-free colorings of graphs, *Theor. Comput. Sci.* 566 (C) (2015) 39–49. doi:10.1016/j.tcs.2014.11.029.
5. I. V. Reddy, Parameterized algorithms for conflict-free colorings of graphs, *Theor. Comput. Sci.* 745 (2018) 53–62. doi:10.1016/j.tcs.2018.05.025.
6. H. L. Bodlaender, S. Kolay, A. Pieterse, Parameterized complexity of conflict-free graph coloring, in: Proceedings of the 16th Intl. Symposium on Algorithms and Data Structures, WADS, 2019, pp. 168–180. doi:10.1007/978-3-030-24766-9_13.

7. A. Agrawal, P. Ashok, M. M. Reddy, S. Saurabh, D. Yadav, FPT algorithms for conflict-free coloring of graphs and chromatic terrain guarding, CoRR abs/1905.01822. arXiv:1905.01822.
8. C. Keller, S. Smorodinsky, Conflict-free coloring of intersection graphs of geometric objects, *Discrete & Computational Geometry* doi:10.1007/s00454-019-00097-8.
9. S. P. Fekete, P. Keldenich, Conflict-free coloring of intersection graphs, *International Journal of Computational Geometry & Applications* 28 (03) (2018) 289–307. doi:10.1142/S0218195918500085.
10. K. Chen, A. Fiat, H. Kaplan, M. Levy, J. Matoušek, E. Mossel, J. Pach, M. Sharir, S. Smorodinsky, U. Wagner, E. Welzl, Online conflict-free coloring for intervals, *SIAM J. Comput.* 36 (5) (2006) 1342–1359. doi:10.1137/S0097539704446682.
11. J. Pach, G. Tardos, Conflict-free colourings of graphs and hypergraphs, *Combinatorics, Probability and Computing* 18 (5) (2009) 819–834. doi:10.1017/S0963548309990290.
12. P. Cheilaris, Conflict-free coloring, Ph.D. thesis, New York, NY, USA (2009).
13. M. Cygan, F. V. Fomin, Ł. Kowalik, D. Lokshtanov, D. Marx, M. Pilipczuk, M. Pilipczuk, S. Saurabh, *Parameterized Algorithms*, 1st Edition, Springer Publishing Company, Incorporated, 2016.
14. N. Robertson, D. Sanders, P. Seymour, R. Thomas, The four-colour theorem, *J. Comb. Theory Ser. B* 70 (1) (1997) 2–44. doi:10.1006/jctb.1997.1750.
15. R. Diestel, *Graph Theory*, Springer-Verlag Heidelberg, 2005.
16. Y. Aubry, J.-C. Godin, O. Togni, Free choosability of outerplanar graphs, *Graphs and Combinatorics* 32 (3) (2016) 851–859. doi:10.1007/s00373-015-1625-3.
17. S. L. Mitchell, Linear algorithms to recognize outerplanar and maximal outerplanar graphs, *Information Processing Letters* 9 (5) (1979) 229 – 232. doi:10.1016/0020-0190(79)90075-9.

A Proofs of Cases 2 and 3 of Theorem 7

A.1 Proof of Case 2

X is a bag that introduces one vertex v_k and $2k - 1 \in \{U(x) : x \in X'\}$. This means that $F(X') \setminus \{U(x) : x \in N(v_k) \cap X'\} \neq \emptyset$. In this case, $C(v_k) = U(w) = 2k - 1$, for a vertex w that is needy in X' , chosen according to Rule 1.

If $C(w) \notin \{1, 2, 3, \dots, 2k - 2\}$, then $(S \cup \{w\}) \setminus \{v_k\}$ is an expensive subset of size k in X' , the predecessor of X . This contradicts the choice of X . Hence $C(w) \in \{1, 2, 3, \dots, 2k - 2\}$. By invariant 1, for any $v, v' \in X'$, we have $C(v) \neq C(v')$. We can rule out the colors $\{1, 3, \dots, 2k - 3\}$ since they appear as $C(v_i)$, for $1 \leq i \leq k - 1$. We can also rule out the colors $\{2, 4, \dots, 2r\}$ since the vertices v_i are needy for $1 \leq i \leq r$. Hence $C(w) \in \{2(r + 1), 2(r + 2), \dots, 2k - 2\}$. Let $C(w) = 2j$ for some $r + 1 \leq j \leq k - 1$. Since $U(v_j) = 2j$, we have $|\{x : x \in X', C(w) = U(x)\}| \geq 1$.

Without loss of generality, let $\{2, 4, \dots, 2\ell\} \subseteq \{U(x) : x \in N(v_k) \cap X'\}$ and $\{2(\ell + 1), \dots, 2r\} \cap \{U(x) : x \in N(v_k) \cap X'\} = \emptyset$ for some $0 \leq \ell \leq r$. Clearly, we cannot choose $C(v_k)$ from $\{2, 4, \dots, 2\ell\}$. Let us try to understand why $2k - 1$ was chosen as $C(v_k)$ over elements of $\{2(\ell + 1), \dots, 2r\}$. We have two subcases.

- $2k - 1 \in F_1(X')$. Then w is the lone vertex in X' such that $U(w) = 2k - 1$. Without loss of generality, let colors $2(\ell + 1), \dots, 2\ell' \in F_1(X')$ and let $2(\ell' + 1), \dots, 2r \in F_{>1}(X')$ for some $\ell < \ell' \leq r$.
As per Rule 1, $2k - 1$ was chosen as a color $c \in F_1(X') \setminus \{U(x) : x \in N(v_k) \cap X'\}$ that minimizes $|\{x : x \in X', C(U^{-1}(c)) = U(x)\}|$. Since $|\{x : x \in X', C(w) = U(x)\}| \geq 1$, for each $\ell + 1 \leq i \leq \ell'$, we have $|\{x : x \in X', C(U^{-1}(2i)) = U(x)\}| = |\{x : x \in X', C(v_i) = U(x)\}| = |\{x : x \in X', 2i - 1 = U(x)\}| \geq 1$. So there exists a set $W' = \{w_{\ell+1}, \dots, w_{\ell'}\}$ (disjoint from S) such that $U(w_i) = 2i - 1$, for each $\ell + 1 \leq i \leq \ell'$.
Since $2(\ell' + 1), \dots, 2r \in F_{>1}(X')$, we have a set $W'' = \{w_{\ell'+1}, \dots, w_r\}$ (disjoint from S) such that $U(w_i) = U(v_i) = 2i$, for each $\ell' + 1 \leq i \leq r$. Thus we have $W = W' \cup W''$ such that $|W| = r - \ell$ that is disjoint from S .
- $2k - 1 \in F_{>1}(X')$. Since a member of $F_{>1}(X')$ was chosen, it follows that $F_1(X') \setminus \{U(x) : x \in N(v_k) \cap X'\} = \emptyset$. Hence $\{2(\ell + 1), \dots, 2r\} \subseteq F_{>1}(X')$. So we have a set $W = \{w_{\ell+1}, \dots, w_r\}$ (disjoint from S) such that $U(w_i) = U(v_i) = 2i$, for each $\ell + 1 \leq i \leq r$. Thus we have W with $|W| = r - \ell$ that is disjoint from S .

If $\ell = 0$, then $|W| = r$, giving us $|X| \geq |S| + |W \cup Z| \geq 3k/2$. In what follows, we will assume $\ell \geq 1$. That is, v_k has at least one needy neighbor. Recall that $\{2, 4, \dots, 2\ell\} \subseteq F(X') \cap \{U(x) : x \in N(v_k) \cap X'\}$. For $1 \leq i \leq \ell$, let $v'_i \in N(v_k) \cap X'$ such that³ $U(v'_i) = 2i$.

Now let us see how $U(v_k)$ was assigned as $2k$. By Rule 2, $U(v_k)$ is set to $C(y)$ such that y is a needy neighbor that minimizes $|\{x : x \in X', U(y) = U(x)\}|$. So $C(y) = 2k$. If $U(y) \notin \{1, 2, \dots, 2k - 2\}$, then $(S \cup \{y\}) \setminus \{v_k\}$ is a k -expensive subset in X' , contradicting the choice of X . So $U(y) \in \{1, 2, 3, \dots, 2k - 2\}$. Since y is needy, as in Case 1, we can rule out $\{1, 3, \dots, 2k - 3\} \cup \{2(r + 1), 2(r + 2), \dots, 2k - 2\}$. So $U(y) \in \{2, 4, 6, \dots, 2r\}$.

Let $U(y) = 2j'$, where $1 \leq j' \leq r$. Since $U(v_{j'}) = 2j'$ as well, we have $|\{x : x \in X', U(y) = U(x)\}| \geq 2$. All of v'_1, \dots, v'_ℓ are needy and neighbors to v_k . Since y was chosen over these vertices, it follows that there exists a set of vertices $Y = \{y_1, \dots, y_\ell\}$, disjoint from S such that $U(y_i) = U(v'_i) = 2i$, for $1 \leq i \leq \ell$.

The sets W and Y are disjoint, but need not be disjoint from Z . Since $|W \cup Y| = r$ and $|Z| = k - r$, we have $|W \cup Y \cup Z| \geq k/2$. Since W, Y, Z are all disjoint from S , we have that $|X| \geq |S| + |W \cup Y \cup Z| \geq 3k/2$. \square

A.2 Proof of Case 3

X is a special bag that introduces v_k and \widehat{v}_k . If $F(X') = \emptyset$, then none of the $k - 1$ vertices in $S \cap X'$ are needy in X' . Hence $|X'| \geq 2(k - 1)$. This implies that $|X| \geq 2(k - 1) + 2 = 2k$ and we are done.

Else, $|F(X')| \geq 1$. Let us first note that since S is an expensive subset, so is $S \cup \{\widehat{v}_k\} \setminus \{v_k\}$. Since $|F(X')| \geq 1$, at least one of $C(v_k)$ or $C(\widehat{v}_k)$ will be chosen from $F(X')$. Without loss of generality, let v_k be a vertex such that

³ The vertices v'_i may or may not be the same as v_i .

$C(v_k) \in F(X')$. Let $C(v_k) = U(w) = 2k - 1$, where w is a needy vertex in X' , chosen according to Rule 1.

If $C(w) \notin \{1, 2, 3, \dots, 2k - 2\}$, then $(S \cup \{w\}) \setminus \{v_k\}$ is an expensive subset of size k in X' , the predecessor of X . This contradicts the choice of X . Hence $C(w) \in \{1, 2, 3, \dots, 2k - 2\}$. By invariant 1, for any $v, v' \in X'$, we have $C(v) \neq C(v')$. We can rule out the colors $\{1, 3, \dots, 2k - 3\}$ since they appear as $C(v_i)$, for $1 \leq i \leq k - 1$. We can also rule out the colors $\{2, 4, \dots, 2r\}$ since the vertices v_i are needy for $1 \leq i \leq r$. Hence $C(w) \in \{2(r + 1), 2(r + 2), \dots, 2k - 2\}$. Let $C(w) = 2j$ for some $r + 1 \leq j \leq k - 1$. Since $U(v_j) = 2j$, we have $|\{x : x \in X', C(w) = U(x)\}| \geq 1$.

Since $\{2, 4, \dots, 2r\} \subseteq F(X')$, let us see why $2k - 1$ was chosen as $C(v_k)$ over these colors. We have two subcases.

- $2k - 1 \in F_1(X')$. Then w is the lone vertex in X' such that $U(w) = 2k - 1$. Without loss of generality, let colors $2, 4, \dots, 2\ell' \in F_1(X')$ and let $2(\ell' + 1), \dots, 2r \in F_{>1}(X')$ for some $0 < \ell' \leq r$. As per Rule 1, $2k - 1$ was chosen as a color $c \in F_1(X')$ that minimizes $|\{x : x \in X', C(U^{-1}(c)) = U(x)\}|$. Since $|\{x : x \in X', C(w) = U(x)\}| \geq 1$, for each $1 \leq i \leq \ell'$, we have $|\{x : x \in X', C(U^{-1}(2i)) = U(x)\}| = |\{x : x \in X', C(v_i) = U(x)\}| = |\{x : x \in X', 2i - 1 = U(x)\}| \geq 1$. So there exists a set $W' = \{w_1, \dots, w_{\ell'}\}$ (disjoint from S) such that $U(w_i) = 2i - 1$, for each $1 \leq i \leq \ell'$. Since $2(\ell' + 1), \dots, 2r \in F_{>1}(X')$, we have a set $W'' = \{w_{\ell'+1}, \dots, w_r\}$ (disjoint from S) such that $U(w_i) = U(v_i) = 2i$, for each $\ell' + 1 \leq i \leq r$. Thus we have $W = W' \cup W''$ such that $|W| = r$ that is disjoint from S .
- $2k - 1 \in F_{>1}(X')$. Since a member of $F_{>1}(X')$ was chosen, it follows that $F_1(X') = \emptyset$. Hence $\{2, 4, \dots, 2r\} \subseteq F_{>1}(X')$. So we have a set $W = \{w_1, \dots, w_r\}$ (disjoint from S) such that $U(w_i) = U(v_i) = 2i$, for each $1 \leq i \leq r$. Thus we have W with $|W| = r$ that is disjoint from S .

In either case, we have W that is disjoint from S and $|W| = r$. Recall that we also have Z disjoint from S , such that $|Z| = k - r$. Thus we get that $|X| \geq |S| + |W \cup Z| \geq 3k/2$. \square

B Proof of Theorem 14

We CFCN color the type graph H using $\chi_{CN}(H)$ colors. Let $C_H : V_H \rightarrow [\chi_{CN}(H)]$ be that coloring and $U_H : V_H \rightarrow [\chi_{CN}(H)]$ be the corresponding assignment of unique colors. Now, we derive a coloring $C : V(G) \rightarrow \{0, 1, 2, \dots, s, s + 1, s + 2\}$ from C_H , with $s = \chi_{CN}(H) + \frac{\text{ind}(G)}{3}$. Also we identify a unique color in the neighborhood of each vertex, denoted $\bar{U} : V(G) \rightarrow \{0, 1, 2, \dots, s, s + 1, s + 2\}$. Let V_1, V_2, \dots, V_t be the type partition of V . We assign colors to the vertices as follows: For each V_i , choose a representative vertex $r_i \in V_i$ and assign $C(r_i) = C_H(i)$. For each V_i , for all vertices $x \in V_i \setminus \{r_i\}$, we assign $C(x) = 0$. We make the below observations.

- Each of the representative vertices r_i has a uniquely colored neighbor, as C_H is a CFCN coloring of H .
- If V_i is a clique, let r_j be the uniquely colored neighbor of r_i (note that r_j can be r_i itself). For each $x \in V_i$, r_j serves as the uniquely colored neighbor.
- If V_i is an independent set such that $C_H(i) \neq U_H(i)$, the uniquely colored neighbor of r_i is the uniquely colored neighbor for all vertices in V_i .

What remains to be handled are the independent sets V_i , such that $C_H(i) = U_H(i)$. We call these type sets V_i (independent sets) as the *bad sets*. We do not consider singleton V_i 's as bad sets. Also, all the representative vertices r_i see a uniquely colored neighbor, regardless of whether V_i is bad or not. Once a bad set V is fixed, we no longer call it a bad set.

Reduction of bad sets: We process the bad sets in iterations. We require at most $\text{ind}(G)/3$ iterations. In iteration ℓ , where $1 \leq \ell \leq \text{ind}(G)/3$, if there exists a bad set V_i , which has at least two neighboring bad sets, we do the following. We call V_i the *lead set* in this iteration. Let the neighboring bad sets of V_i be $V_{i_1}, V_{i_2}, \dots, V_{i_m}$, where $m \geq 2$. Choose a vertex $v_i \in V_i \setminus \{r_i\}$ and reassign $C(v_i) = \chi_{CN}(H) + \ell$. We call v_i as the *lead representative* of this iteration. For each vertex $x \in V_i \setminus \{r_i, v_i\}$, reassign $C(x) = \chi_{CN}(H) + \ell + 1$. Each vertex in V_i serves as its own the uniquely colored neighbor. For each vertex in V_{i_p} , the vertex $v_i \in V_i$ serves as the uniquely colored neighbor.

Let us see why the reduction operation fixes all the lead sets and all their neighboring bad sets. None of the sets chosen as the lead sets in two different iterations are adjacent, else they could have been considered in the same iteration. For bad sets that neighbors multiple lead sets, the uniquely colored neighbor is provided by the lead representative that was considered earliest in the reduction operation.

In each of the above iterations, at least 3 bad sets are colored. Hence it suffices to have $\text{ind}(G)/3$ iterations. The number of colors required are $\text{ind}(G)/3 + 1$. After the reduction operations, we are left with the bad sets which have at most one bad set as neighbor. We handle them as follows.

- **Case 1: Bad sets V_i and V_j which are neighbor, each of which is not neighbors to any other bad sets.**

We note that the color $\chi_{CN}(H) + \frac{\text{ind}(G)}{3} + 1 = s + 1$ is possibly used only in the last iteration of the reduction operation, but does not serve as a unique color for any of the vertices in the bad sets of that iteration.

We use the colors $s + 1$ and $s + 2$ for coloring the bad sets V_i and V_j . Choose two vertices $x_i \in V_i \setminus \{r_i\}$ and $x_j \in V_j \setminus \{r_j\}$. Reassign $C(x_i) = s + 1$ and $C(x_j) = s + 2$. These vertices x_i and x_j serve as uniquely colored neighbors for the vertices in V_j and V_i respectively.

- **Case 2: A bad set V_i that has no bad set as neighbor.**

Reassign $C(v) = s + 2$, for all vertices $v \in V_i \setminus \{r_i\}$. All the vertices in V_i serve as their own uniquely colored neighbors.

The above coloring is a CFCN coloring. We use $\chi_{CN}(H)$ colors to color the representative vertices of each V_i . and $\text{ind}(G)/3 + 1$ colors in the reduction

operation. Taking the colors $\{0, s + 2\}$ into account, the total number of colors used is $\chi_{CN}(H) + \text{ind}(G)/3 + 3$. \square

C Proof of Theorem 17

Let $\text{dc}(G) = d$. That is, there is a set $X \subseteq V$, with $|X| = d$ such that $G[V \setminus X]$ is a disjoint union of cliques. Let $X = \{v_1, v_2, \dots, v_d\}$ and $Y = \{v_i \in X : \deg_X(v_i) \geq 1\}$.

We have three cases and in each case, we explain how to get CFCN coloring. Cases 1 and 2 use $d + 1$ colors and case 3 uses 3 colors.

1. There is a clique $K' \subseteq G[V \setminus X]$, with $u \in K'$, such that $|N(u) \cap (X \setminus Y)| \geq 2$.
 - Without loss of generality, let $v_{i_1}, v_{i_2}, \dots, v_{i_m} \in N(u) \cap (X \setminus Y)$, where $m \geq 2$.
 - Assign $C(u) = i_1$ and $C(v_{i_\ell}) = d + 1$, for all $1 \leq \ell \leq m$.
Note that the color i_2 is not assigned and will be used for future coloring.
 - For each of the uncolored vertices $v_i \in X \setminus Y$, $C(v_i) = i$.
 - For each of the cliques $K \subseteq G[V \setminus X]$,
 - If K has a colored vertex, color the remaining vertices with $d + 1$.
 - Else, choose a vertex in K and assign the color i_2 . Color the remaining vertices with $d + 1$.

For the vertices $v_{i_\ell} \in N(u) \cap (X \setminus Y)$, where $1 \leq \ell \leq m$, the vertex u is the uniquely colored neighbor. For all the other vertices $v_i \in X$, v_i itself is the uniquely colored neighbor. For all the vertices in cliques $K \subseteq G[V \setminus X]$, the vertex colored i_1 or i_2 will serve as the uniquely colored neighbor.

2. $Y \neq \emptyset$. That is, there exists two vertices $v_i, v_j \in X$ such that $\{v_i, v_j\} \in E(G)$.
 - Assign $C(v_i) = i$ and $C(v_j) = d + 1$.
Note that the color j is not used and will be used for future coloring.
 - For each of the uncolored vertices $v_k \in X$, assign $C(v_k) = k$.
 - For each of the cliques K , choose a vertex and assign the color j . Color the rest of the vertices with $d + 1$.

Each vertex in $X \setminus \{v_j\}$ serves as its own uniquely colored neighbor. For the vertex v_j , the uniquely colored neighbor is v_i . For each clique K , the vertex colored j is the uniquely colored neighbor for all the vertices in K .

3. Else, (i) X is an independent set and (ii) for each clique K , and for all $w \in K$, we have $|N(w) \cap X| \leq 1$.
 - For each clique K , choose a vertex and assign the color 1 and color the remaining vertices with 2.
 - For all vertices $x \in X$, assign $C(x) = 3$.

Note that this is a CFCN 3-coloring of G . Each vertex in X serves as its own uniquely colored neighbor. For each clique K , the vertex colored 1 acts as a uniquely colored neighbor for all the vertices in K . \square

D Proof of Theorem 22

In this section, whenever we refer to an outerplanar graph G , we will also be implicitly referring to a planar drawing of G with all the vertices appearing in the outer face. We will abuse language and say “faces of G ” when we want to refer to faces of the above planar drawing.

Theorem 22 is proved using a two-level induction process. The first level is using a *block decomposition* of the graph. Any connected graph can be viewed as a tree of its constituent blocks. We color the blocks in order so that when we color a block, at most one of its vertices is previously colored. Each block is colored without affecting the color of the already colored vertex. The second level of the induction is required for coloring each of the blocks. We use ear decomposition on each block and color the faces of the block in sequence. However, the proof is quite technical and involves several cases of analysis at each step.

We summarize the relevant aspects of block decomposition below. The reader is referred to a standard textbook in graph theory [15] for more details on this.

- A *block* is a maximal connected subgraph without a cut vertex.
- Blocks of a connected graph are either maximal 2-connected subgraphs, or edges (the edges which form a block will be bridges).
- Two distinct blocks overlap in at most one vertex, which is a cut vertex.
- Any connected graph can be viewed as tree of its constituent blocks.

In the following discussion, we explain how to construct a coloring $C : V(G) \rightarrow \{1, 2, 3, 4\}$ for an outerplanar graph G . At any intermediate stage, the coloring C will satisfy⁴ the following invariants:

Invariants of C

- Every vertex v that has already been assigned a color $C(v)$ has a neighbor w , such that $C(w) \neq C(x)$, for all $x \in N(v) \setminus \{w\}$. For v , the function $U : V(G) \rightarrow \{1, 2, 3, 4\}$ denotes the color of w , its uniquely colored neighbor.
- $\forall v \in V(G), C(v) \neq U(v)$.
- $\forall \{v, w\} \in E(G), C(v) \neq C(w)$ and $|\{C(v), U(v), C(w), U(w)\}| = 3$. (\star)

Theorem 22 is proved by using an induction on the block decomposition of the graph G and the below results.

Lemma 23. *If G is a 2-connected outerplanar graph such that all its inner faces contain exactly 5 vertices, then G has a CFON coloring using 3 colors.*

Theorem 24. *Let G be an outerplanar graph.*

⁴ The condition marked \star is violated in a few cases. In the exceptional cases where it is violated, we shall explain how the cases are handled.

1. If B is a block of G that is either a bridge, or contains an inner face F with $|V(F)| \neq 5$, then B has a CFON coloring using at most 4 colors.
2. If B is a block of G , with exactly one vertex v precolored with color $C(v)$ and unique color $U(v)$, then the rest of B has a CFON coloring using at most 4 colors, while retaining $C(v)$ and $U(v)$.

Proof (Proof of Theorem 22). Let G be an outerplanar graph. We apply block decomposition on G which results in blocks that are either maximal 2-connected subgraphs or single edges.

If G is 2-connected and all its inner faces have exactly 5 vertices, then by Lemma 23, G has a CFON coloring using 3 colors.

If G does not fit the above description, then G has a block B such that either B is an edge, or B has an inner face F with $|V(F)| \neq 5$. In this case, by Theorem 24.1, B has a CFON coloring using at most 4 colors.

Viewing G as a tree of its blocks, we can start coloring blocks that are adjacent to blocks that are already colored. Suppose the block B is already colored, and let B' be a block adjacent to B . Let x be the cut-vertex between the blocks B and B' . We use Theorem 24.2 to obtain a CFON coloring of B' using at most 4 colors, while retaining $C(x)$ and $U(x)$. \square

We now proceed towards proving Lemma 23 and Theorem 24. Lemma 23 and Theorem 24 discusses the coloring of blocks, which is accomplished by means of induction on the faces of the blocks. Towards this end, we use the following fact about ear decomposition of 2-connected outerplanar graphs. For a proof of the below lemma, we refer the reader to [16] where this is stated as Observation 2.

Lemma 25 (Ear Decomposition). *Let B be a 2-connected block in an outerplanar graph. Then B has an ear decomposition $F_0, P_1, P_2, \dots, P_q$ satisfying the following:*

- F_0 is an arbitrarily chosen inner face of B .
- Every P_i is a path with end points v, w such that $\{v, w\}$ is an edge in $F_0 \cup \bigcup_{1 \leq j < i} P_j$. Thus P_i together with the edge $\{v, w\}$ forms a face of B .

We are now ready to prove Lemma 23.

Proof. (Proof of Lemma 23) Since G is 2-connected, the entire graph forms a single block. Let F_0, P_1, \dots, P_q be an ear decomposition of G . Recall that all the faces have exactly five vertices. Let $F_0 = v_1 - v_2 - v_3 - v_4 - v_5 - v_1$. We assign⁵ the following colors to the vertices in F_0 : $C(v_1) = 1, C(v_2) = 1, C(v_3) = 2, C(v_4) = 2, C(v_5) = 3$. We also have $U(v_1) = 3, U(v_2) = 2, U(v_3) = 1, U(v_4) = 3, U(v_5) = 1$.

Let P_i be any subsequent face $P_i = w_1 - w_2 - w_3 - w_4 - w_5 - w_1$ with $\{w_1, w_2\}$ being the pre-existing edge in $F_0 \cup \bigcup_{1 \leq j < i} P_j$. Depending on the values already assigned to $C(w_1), U(w_1), C(w_2), U(w_2)$, we assign the colors to w_3, w_4 and w_5 .

⁵ The coloring assigned in this proof does not satisfy the condition marked \star . However, this is not an issue since we are coloring the whole of G in this lemma.

We always ensure that $C(v) \neq U(v)$ for all vertices v . We note that the values $C(w_1), U(w_1), C(w_2), U(w_2)$ can take only the four below combinations, w.l.o.g. We explain the coloring for the rest of P_i in each of these cases.

1. $C(w_1) = C(w_2)$ and $|\{C(w_1), U(w_1), U(w_2)\}| = 3$. W.l.o.g., let $C(w_1) = 1, U(w_1) = 2, C(w_2) = 1, U(w_2) = 3$. Assign $C(w_3) = 2, C(w_4) = 2, C(w_5) = 3$ and $U(w_3) = 1, U(w_4) = 3, U(w_5) = 1$.
2. $C(w_1) \neq C(w_2), U(w_1) \neq U(w_2)$, and $|\{C(w_1), C(w_2), U(w_1), U(w_2)\}| = 3$. Either w_1 serves as the uniquely colored neighbor of w_2 or vice versa. W.l.o.g., let $C(w_1) = 1, U(w_1) = 2, C(w_2) = 2, U(w_2) = 3$. Assign $C(w_3) = 1, C(w_4) = 3, C(w_5) = 3$ and $U(w_3) = 2, U(w_4) = 1, U(w_5) = 1$.
3. $C(w_1) = U(w_2)$ and $C(w_2) = U(w_1)$. W.l.o.g., let $C(w_1) = 1, U(w_1) = 2, C(w_2) = 2, U(w_2) = 1$. Assign $C(w_3) = 2, C(w_4) = 3, C(w_5) = 1$ and $U(w_3) = 3, U(w_4) = 2, U(w_5) = 3$.
4. $C(w_1) = C(w_2)$ and $U(w_1) = U(w_2)$. W.l.o.g., let $C(w_1) = C(w_2) = 1, U(w_1) = U(w_2) = 2$. Assign $C(w_3) = 1, C(w_4) = 2, C(w_5) = 3$ and $U(w_3) = 2, U(w_4) = 3, U(w_5) = 1$.
5. The case $U(w_1) = U(w_2)$ and $|\{U(w_1), C(w_1), C(w_2)\}| = 3$ does not arise in the above colorings.

□

At this point, to complete the proof of Theorem 22 we need to prove Theorem 24. We now state a few results that would help us towards this end.

Lemma 26. *An uncolored face F , such that $|V(F)| \neq 5$, can be CFON colored using 4 colors satisfying the invariants.*

Proof. Let $F = v_1 - v_2 - v_3 - \dots - v_{k-1} - v_k - v_1$ be a face with $|V(F)| = k$, $k \neq 5$. We assign $C(v_1) = 1, C(v_2) = 2, C(v_3) = 3$ and for the remaining vertices (if any), we set $C(v_i) = C(v_{i-3})$. In order to satisfy the invariants, we need to make the following changes:

- $k \equiv 0 \pmod{3}$. No change is necessary.
- $k \equiv 1 \pmod{3}$. Reassign $C(v_k) = 4$.
- $k \equiv 2 \pmod{3}$. Reassign $C(v_{k-3}) = 4, C(v_{k-2}) = 2, C(v_{k-1}) = 3, C(v_k) = 4$. Notice that this coloring does not satisfy the invariants if $k = 5$. However, the smallest k that we consider in this case is $k = 8$.

In each of the above cases the unique color for each vertex v_i is provided by its cyclical successor i.e., $U(v_i) = C(v_{i+1})$. □

Lemma 27. *Let F be a face (cycle) in G with one vertex v such that $C(v)$ and $U(v)$ are already assigned, with $C(v) \neq U(v)$. Then the rest of F can be CFON colored using at most 4 colors, while retaining $C(v)$ and $U(v)$, and satisfying the invariants.*

Proof. Let v_1 be the colored vertex in the cycle F . We may assume w.l.o.g. that $C(v_1) = 1$ and $U(v_1) = 2$. Now, we extend C to the remainder of F .

- $|V(F)| = 3$ with $F = v_1 - v_2 - v_3 - v_1$.
We assign: $C(v_2) = 3$, $C(v_3) = 4$ and $U(v_2) = 1$, $U(v_3) = 1$.
- $|V(F)| \geq 4$ with $F = v_1 - v_2 - v_3 - \dots - v_{k-1} - v_k - v_1$.
We first assign: $C(v_2) = 3$ and $C(v_3) = 2$. For the remaining vertices v_i , we set $C(v_i) = C(v_{i-3})$ for $4 \leq i \leq k$. However, we need to make some changes to this in order to satisfy the invariants. We have the following subcases:
 - $k \equiv 0$ or $1 \pmod{3}$. Reassign $C(v_k) = 4$.
 - $k \equiv 2 \pmod{3}$. Reassign $C(v_{k-1}) = 4$.
 In each of the above cases the unique color for each vertex v_i is provided by its cyclical successor i.e., $U(v_i) = C(v_{i+1})$. Observe that $U(v_1)$ is left unchanged, by ensuring v_2 and v_k , the neighbors of v_1 , are not assigned the color $U(v_1)$.

□

Lemma 28. *Let F be a face with $|V(F)| \geq 4$ with such that the edge $\{v_1, v_2\} \in E(F)$ and v_1 and v_2 already colored such that $C(v_1) = C(v_2)$ and $U(v_1) \neq U(v_2)$. Then the rest of F can be CFON colored using 4 colors satisfying the invariants.*

Proof. W.l.o.g., we may assume $C(v_1) = C(v_2) = 4$, $U(v_1) = 1$ and $U(v_2) = 2$. We have the following cases:

- $|V(F)| = 4$ with $F = v_2 - v_3 - v_4 - v_1 - v_2$. We assign: $C(v_3) = 1$, $C(v_4) = 3$ and $U(v_3) = 4$ and $U(v_4) = 4$.
- If $|V(F)| = 5$ with $F = v_2 - v_3 - v_4 - v_5 - v_1 - v_2$. We assign: $C(v_3) = 1$, $C(v_4) = 2$, $C(v_5) = 3$ and $U(v_3) = 2$, $U(v_4) = 3$ and $U(v_5) = 4$.
- If $|V(F)| \geq 6$ with $F = v_2 - v_3 - \dots - v_{k-1} - v_k - v_1 - v_2$. We assign: $C(v_3) = 3$ and $C(v_4) = 2$. For all $5 \leq i \leq k$, $C(v_i) = C(v_{i-3})$.
 - $k \equiv 0 \pmod{3}$. Reassign $C(v_{k-1}) = 1$.
 - $k \equiv 1 \pmod{3}$. No change is required.
 - $k \equiv 2 \pmod{3}$. Reassign $C(v_{k-1}) = 1$ and $C(v_k) = 2$.
 The unique color of each vertex v_i is provided its cyclical successor i.e., $U(v_i) = C(v_{i+1})$.

□

Lemma 29. *Let P be a path in G whose endpoints are v_1, v_2 . Suppose $\{v_1, v_2\} \in E(G)$ and that v_1, v_2 are already assigned the functions C and U satisfying the invariants. Then the rest of P can be CFON colored using at most 4 colors, while retaining C and U values of the endpoints, and satisfying the invariants.*

Since the proof of the above lemma is a bit long and involved, we first prove Theorem 24 using Lemmas 26, 27 and 29.

Proof (Proof of Theorem 24).

1. If the block is a bridge, say $\{v, w\}$, then we color it $C(v) = 1, C(w) = 2$ with $U(v) = 2, U(w) = 1$. Note that the invariant marked \star is violated in this case. However, this does not cause an issue since this edge is a bridge, and it does not appear in any inner face.

If the block is not a bridge, then by assumption, it contains a face F such that $|V(F)| \neq 5$. By Lemma 26, we have a coloring of F using 4 colors and satisfying the invariants. By the Lemma 25 (Ear Decomposition), the block has an ear decomposition F, P_1, P_2, \dots with F as the starting inner face. Recall that for every path P_i , the end points form an edge in $F_0 \cup \bigcup_{1 \leq j < i} P_j$. We color the paths P_1, P_2, \dots in this order. By Lemma 29, we have a coloring for each of these paths using 4 colors and satisfying the invariants.

2. Let v be the vertex in the block that is already colored. W.l.o.g., we may assume that $C(v) = 1$ and $U(v) = 2$.

If the block is a bridge $\{v, w\}$, we color w with $C(w) = 3$ and set $U(w) = 1$.

If the block is not a bridge, choose an inner face F that contains v . Using Lemma 27, we color the remainder of F using at most 4 colors and satisfying the invariants. The rest of the proof follows from the fact that we have an ear decomposition with F as the starting face, and Lemma 29. This is very similar to the argument in the proof of part 1 of this theorem and hence the details are omitted.

□

In order to complete the proof of Theorem 22, the last remaining piece is the proof of Lemma 29.

Proof (Proof of Lemma 29). Let v_1 and v_2 be the end points of P . We extend the coloring C to the remainder of P . According to the invariants of C , we have only 2 cases possible.

Case 1: $C(v_1) \neq C(v_2), U(v_1) \neq U(v_2)$. W.l.o.g. we may assume $C(v_1) = 1, C(v_2) = 2$ and $U(v_1) = 2, U(v_2) = 3$.

- $|V(P)| = 3, P = v_2 - v_3 - v_1$. Assign $C(v_3) = 4$ with $U(v_3) = 2$.
- $|V(P)| = 4, P = v_2 - v_3 - v_4 - v_1$. Assign $C(v_3) = 4, C(v_4) = 3$ with $U(v_3) = 3, U(v_4) = 1$.
- $|V(P)| \geq 5, P = v_2 - v_3 - \dots - v_{k-1} - v_k - v_1$. We first assign $C(v_3) = 1, C(v_4) = 3, C(v_5) = 4$. For the remaining vertices v_i , we initially assign $C(v_i) = C(v_{i-3})$ for $6 \leq i \leq k$. However, we need to make some changes to satisfy the invariants. We have the following subcases:
 - $k \equiv 0 \pmod{3}$. Reassign $C(v_{k-1}) = 2$ and $C(v_k) = 4$
 - $k \equiv 1 \pmod{3}$. Reassign $C(v_{k-1}) = 2$.
 - $k \equiv 2 \pmod{3}$. No change is necessary.

In each of the above cases the unique color for each vertex v_i is provided by its cyclical successor i.e., $U(v_i) = C(v_{i+1})$.

Case 2: $U(v_1) = U(v_2)$. W.l.o.g., we may assume $C(v_1) = 1, C(v_2) = 2$ and $U(v_1) = U(v_2) = 3$.

- **Case 2(i):** $|V(P)| = 3$ and $P = v_2 - v_3 - v_1$.
 - **Case 2(i)(a):** Vertices v_1 and v_2 are the only neighbors of v_3 . Assign $C(v_3) = 4$ and $U(v_3) = 2$. The invariant marked \star is not satisfied, but that does not matter as v_3 does not participate in any further faces.




Fig. 1. Case 2(i)(d)

- **Case 2(i)(b):** One of the edges $\{v_1, v_3\}$ or $\{v_2, v_3\}$ does not feature in an another face. W.l.o.g., say $\{v_2, v_3\}$ be that edge. Assign $C(v_3) = 4$ with $U(v_3) = 1$. The \star invariant is violated for $\{v_2, v_3\}$ here but it does not affect the further coloring.
 - **Case 2(i)(c):** One of the edges $\{v_1, v_3\}$ or $\{v_2, v_3\}$ features in an uncolored face F such that $|V(F)| \neq 3$. W.l.o.g., say $\{v_2, v_3\}$ is that edge. We assign $C(v_3) = 4$ with $U(v_3) = 1$. Let $|V(F)| = k$ with $F = v_3 - w_1 - w_2 - \dots - w_{k-2} - v_2 - v_3$. We assign $C(w_1) = 3$, $C(w_2) = 1$ and $C(w_3) = 4$ (if w_3 exists). For all $4 \leq i \leq k-2$, $C(w_i) = C(w_{i-3})$. If $k \equiv 0 \pmod{3}$, we reassign $C(w_{k-4}) = 2$, $C(w_{k-3}) = 1$ and $C(w_{k-2}) = 4$. The unique colors U for the vertices are assigned as follows:
 - * For $k = 6$, $U(w_1) = 4, U(w_2) = 3, U(w_3) = 2$ and $U(w_4) = 2$.
 - * For $k \neq 6$, we have for $1 \leq i \leq k-3$, $U(w_i) = C(w_{i+1})$ and $U(w_{k-2}) = C(v_2) = 2$.
 - **Case 2(i)(d):** The only remaining case is when both the edges $\{v_1, v_3\}$ or $\{v_2, v_3\}$ feature in uncolored triangular faces. Let $\{v_1, v_3\}$ form a triangular face with x and $\{v_2, v_3\}$ with y . We have two subcases:
 - * The edge $\{x, v_3\}$ forms a triangular face with another vertex z (see Figure 1). Assign $C(v_3) = 1, C(x) = 2, C(y) = 4, C(z) = 3$ and $U(v_3) = 4, U(x) = 3, U(y) = 2, U(z) = 1$. Some edges violate the invariant marked \star , but these edges are already part of two faces, and hence do not feature in the further coloring.
 - * The edge $\{x, v_3\}$ is not part of a triangular face with another vertex. In this case, we assign $C(v_3) = 4, C(x) = 4, C(y) = 1$ and $U(v_3) = 2, U(x) = 1, U(y) = 2$. Out of the edges that violate the invariant marked \star , the only one that can participate in the further coloring is the edge $\{x, v_3\}$. By assumption, $\{x, v_3\}$ is not part of a triangular face. In Lemma 28, we explain how to color the uncolored face that is $\{x, v_3\}$ may be a part of.
- **Case 2(ii):** $|V(P)| = 4$, $P = v_2 - v_3 - v_4 - v_1$.
- **Case 2(ii)(a):** The edge $\{v_3, v_4\}$ forms a triangular face with a vertex x . We assign $C(v_3) = 1, C(v_4) = 4, C(x) = 3$, with $U(v_3) = 3, U(v_4) = 1, U(x) = 4$.

- **Case 2(ii)(b):** The edge $\{v_3, v_4\}$ is not part of an uncolored triangular face. We assign $C(v_3) = C(v_4) = 4$, with $U(v_3) = 2, U(v_4) = 1$. If the edge $\{v_3, v_4\}$ is part of an uncolored face F , by assumption, we know that $|V(F)| \geq 4$ and hence we can use Lemma 28 to color F satisfying the invariants.
- **Case 2(iii):** $|V(P)| = 5$ with $P = v_2 - v_3 - v_4 - v_5 - v_1$. We assign $C(v_3) = 1, C(v_4) = 3, C(v_5) = 2$, with $U(v_3) = 3, U(v_4) = 2, U(v_5) = 1$.
- **Case 2(iv):** $|V(P)| \geq 6$, with $P = v_2 - v_3 - \dots - v_{k-2} - v_{k-1} - v_k - v_1$. We first assign $C(v_3) = 4$ and $C(v_4) = 3$. For $5 \leq i \leq k$, assign $C(v_i) = C(v_{i-3})$. If $k \equiv 1 \pmod{3}$, then reassign $C(v_{k-2}) = 1$ and $C(v_k) = 2$. For each vertex v_i , the unique color is provided by its cyclical successor i.e., $U(v_i) = C(v_{i+1})$.

□

Algorithmic Note: The steps in the proof of Theorem 22 leads to an algorithm. Block decomposition, outerplanarity testing and embedding outerplanar graphs [17] can all be done in linear time, i.e., $O(|V(G)|)$. Thus we have an $O(|V(G)|)$ time algorithm, that given an outerplanar graph G , determines a CFON coloring for G that uses four colors.